

REFERENCES

- Arvidsson, P., M. van Boekel, K. Skog, A. Solyakov, and M. Jagerstad. 1999. Formation of heterocyclic amines in a meat juice model system. *Journal of Food Science*. **64**:216-221.
- Balogh, Z., J.I. Gray, E.A. Gomaa, and A.M. Booren. 2000. Formation and inhibition of heterocyclic aromatic amines in fried ground beef patties. *Food Chem Toxicol.* **38**:395-401.
- Bartle, K.D. 1993. Introduction to the theory of chromatographic separations. In Baugh, P.J. (eds.), *Gas Chromatography A practical Approach.* (vol.I), pp.1-14. New York: Oxford University Press.
- Chen, B.H. and D.J. Yang. 1998. An improved analytical method for determination of heterocyclic amines in chicken legs. *Chromatographia*. **48**:223-230.
- DeBruin, L.S., P.A. Martos, and P.D. Josephy. 2001. Detection of PhIP (2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine) in the milk of healthy women. *Chem Res Toxicol.* **14**:1523-1528.
- <http://www.ipmofalaska.com/files/DE.html>. Diatomaceous Earth and Silica Aerogel.
- EURACHEM guide, The fitness for Purpose of Analytical Methods: A Laboratory Guide to Method Validation and Related Topics, update on December, 1998. <http://www.fasor.com>

- Fan, L., Schut, H.A. and Snyderwine, E.G. 1995. Cytotoxicity, DNA adduct formation and DNA repair induced by 2-hydroxyamino-3-methyl imidazo [4,5-f]quinoline and 2-hydroxyamino-1-methyl-6-phenyl imidazo[4,5-b]pyridine in cultured human mammary epithelial cells. *Carcinogenesis.* **16**:775-779.
- Galceran, M.T., Pais,P. and Puignou, L. 1993. High-Performance Liquid-Chromatographic Determination of 10 Heterocyclic Aromatic-Amines with Electrochemical Detection. *J. Chromatography A.* **655**:101-110.
- Grob, R.L. 1985. Modern Practice of Gas Chromatography (2nd ed.), pp. 50-114. United States of America: John Wiley&Sons,.Inc.
- Gross, G.A. and A. Grüter. 1992. Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products. *J Chromatogr.* **592**: 271-278.
- Gross, G.A., R.J. Turesky, L.B. Fay, W.G. Stillwell, P.L. Skipper, and S.R. Tannenbaum. 1993. Heterocyclic aromatic amine formation in grilled bacon, beef and fish and in grill scrapings. *Carcinogenesis.* **14**:2313-2318.
- IARC. 1993. Some Naturally Occurring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines, and Mycotoxins. IQ (2-amino-3-methylimidazo[4,5-f]quinoline). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Vol 56. Lyon, France, World Health Organization.

IARC. 1993a. PhIP (2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine).

In IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Some Naturally Occurring Substances - Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins, Vol 56. World Health Organization, International Agency for Research on Cancer, Lyon, France. pp. 229-242.

IARC. 1993b. MeIQ (2-amino-3,4-dimethylimidazo[4,5-f]quinoline). In IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Some Naturally Occurring Substances - Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins, Vol 56. World Health Organization, International Agency for Research on Cancer, Lyon, France. pp. 197-210.

IARC. 1993c. MeIQx (2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline). In IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Some Naturally Occurring Substances - Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins, Vol 56. World Health Organization, International Agency for Research on Cancer, Lyon, France. pp. 211-228.

Jackson, L.S., Hargraves, W.A. Stroup, W.H. and Diachenko, G.W. 1994. Heterocyclic aromatic amine content of selected beef flavors. *Mutat Res.* 320:113-124.

Janoszka, B., Blaszezyk, U., Warzecha, L., Strózyk, M., Damasiewicz heterocyclic aromatic amines (aminoazaarenes) from heat-treated meat samples. *J.Chromatogr A.* 938:155-165.

- Ji, H., Yu, M.C., Stillwell, W.G., Skipper, P.L., Ross, R.K., Henderson, B.E. and Tannenbaum, S.R. 1994. Urinary excretion of 2-amino-3,8-dimethylimidazo-[4,5-f]quinoxaline in white, black, and Asian men in Los Angeles County. *Cancer Epidemiol Biomarkers Prev* **3**:407-411.
- Kataoka, H. 1996. Review Derivatization reactions for the determination of amines by gas chromatography and their applications in environmental analysis. *J. Chromatogr. A*, **733**:19-34
- Kataoka, H. and Kijima, K. 1997. Analysis of heterocyclic amines as their N-dimethylaminomethylene derivatives by gas chromatography with nitrogen-phosphorus selective detection. *J. Chromatogr. A*, **767**:187-194.
- Knize, M.G., Sinha, R., Rothman, N., Brown, E.D., Salmon, C.P., Levander, O.A., Cunningham, P.L. and Felton, J.S. 1995. Heterocyclic amine content in fast-food meat products. *Food Chem Toxicol* **33**:545-551.
- Keating, G.A., Layton, D.W. and Felton, J.S. 1999. Factors determining dietary intakes of heterocyclic amines in cooked foods. *Mutation Research* **443**:149-156.
- Leong-Morgenthaler, P.M., Op, H., Jaccaud, V. E. and Turesky, R.J. 1998. Mutagenicity of 2-amino-3-methylimidazo[4,5-f]quinoline in human lymphoblastoid cells. *Carcinogenesis* **19**:1749-1754.
- Lynch, A.M., Murray, S., Gooderham, N.J. and Boobis, A.R. 1995. Exposure to and activation of dietary heterocyclic amines in humans. *Crit Rev Oncol Hematol*. **21**:19-31.

- Manabe, S., Suzuki, H. Wada, O. and Ueki, A. 1993. Detection of the carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in beer and wine. *Carcinogenesis*. **14**:899-901.
- Nutrition Research Nation Cattlemen's Beef Association. Mutagens/Carcinogens And Meat. www.beefnutrition.org.
- Ohe, T. 1997. Quantification of mutagenic/carcinogenic heterocyclic amines, MeIQx, Trp-P-1, Trp-P-2 and PhIP, contributing highly to genotoxicity of river water. *Mutat Res.* **393**:73-79.
- Probst, M.R., Blum, M., Fasshauer, I., D'Orazio, D., Meyer, U.A. and Wild, D. 1992. The role of the human acetylation polymorphism in metabolic activation of the food carcinogen 2-aminomethylimidazo[4,5-f]quinoline (IQ). *Carcinogenesis*. **13**:1713-1717.
- Pais, P. and Knize, M.G. 2000. Review Chromatographic and related techniques for the determination of aromatic heterocyclic amines in foods. *J.Chromatogr B*. **747**: 139-169.
- Pais, P., Salmon, C.P. Knize, M.G. and Felton, J.S. 1999. Formation of mutagenic/carcinogenic heterocyclic amines in dry-heated model systems, meats, and meat drippings. *J Agric Food Chem.* **47**:1098-1108.
- Perkin Elmer. 1995. Gas chromatography AutoSystem XL GC User's Manual. (vol. 1) United States of America.
- Raymond, P.W. 1998. Introduction to Analytical Gas Chromatography (2nd ed.), (vol 76). pp.184-185. New Jersey: Cherry Hill.

- Reistad, R., Nyholm, S.H., Haug L.S., Becher, G. and Alexander, J. 1999. 2-amino-1-methyl-6-phenylimidazo(4,5-*b*)pyridine (PhIP) in human hair as biomarker for dietary exposure. *Biomarkers*. **4**:263-271.
- Richling, E., Decker, C., Haring, D., Herderich, M. and Schreier, P. 1997. Analysis of heterocyclic aromatic amines in wine by high-performance liquid chromatography-electrospray tandem mass spectrometry. *J. Chromatog A*. **791** : 71-77.
- Sinha, R., Rothman, N., Brown, E.D., Mark, S.D., Hoover, R.N., Caporaso, N.E., Levander, O.A., Knize, M.G., Lang, N.P. and Kadlubar, F.F. 1994. Pan-fried meat containing high levels of heterocyclic aromatic amines but low levels of polycyclic aromatic hydrocarbons induces cytochrome P4501A2 activity in humans. *Cancer Res.* **54**:6154-6159.
- Sinha, R., Rothman, N., Brown, E.D., Salmon, C.P., Knize, M.G., Swanson, C.A., Rossi, S.C., Mark, S.D., Levander, O.A. and Felton, J.S. 1995. High concentrations of the carcinogen 2-amino-1-methyl-6-phenylimidazo-[4,5-*b*]pyridine (PhIP) occur in chicken but are dependent on the cooking method. *Cancer Res.* **55**:4516-4519.
- Skog, K., Augustsson, K., Steineck, G., Stenberg, M. and Jagerstad, M. 1997. Polar and non-polar heterocyclic amines in cooked fish and meat products and their corresponding pan residues. *Food Chem Toxicol.* **35**:555-565.
- Skog, K. and Solyakov, A. 2002. Screening for heterocyclic amines in chicken cooked in various ways. *Food and Chemical Toxicology*. **40**:1205-1211.

Sugimura, T. 2002. Food and Cancer. *Toxicology*. **17-21** :181-182.

Technology Planning and Management Corporation. 2002. Report on Carcinogens Background Document for Draft RoC Background Document for Selected Heterocyclic Amines. U.S. Department of Health and Human Services Public Health Service National Toxicology Program Research Triangle Park, NC 27709 13 September 2002, pp.1-245.

Technology Planning and Management Corporation. Draft Report on Carcinogens Background Document for 2-Amino-3-methylimidazo [4,5-f]quinoline (IQ) Meeting of the NTP Board of Scientific Counselors Report on Carcinogens Subcommittee. U.S. Department of Health and Human Services Public Health Service National Toxicology Program Research Triangle Park, NC 27709

Thiébaud, H.P., M.G. Knize, P.A. Kuzmicky, D.P. Hsieh, and J.S. Felton. 1995. Airborne mutagens produced by frying beef, pork and a soy-based food. *Food Chem Toxicol* **33**:821-828.

Toribio, F., Galceran, M.T. and Puignou, L. 2000. Review Separation of heterocyclic amines in food product. *J. Chromatogr B.*, **747** 171-202.

Toribio, F., Moyano, E., Puignou, L., and Galceran, M.T. 2002. Ion-trap tandem mass spectrometry for the determination of heterocyclic amines in food. *J. Chromatogr A.*, **948**:267-281.

Townshend A. 1995. Encyclopedia of Analytical Science (1st ed.), London: Academic Press Limited.

Tripler, A. 1993. Gas chromatographic instrumentation, and considerations. In Baugh, P.J. (eds.), *Gas Cromatography A practical Approach.* (vol.I), pp.35-36. New York: Oxford University Press.