

CONTENTS

	page
CONTENTS	x
LIST OF TABLES	xiv
LIST OF FIGURES	xviii
CHAPTER I : INTRODUCTION AND LITERATURE REVIEW	1
Biological properties of carotenoid	5
Carotenoid in photosynthetic bacteria	10
Biosynthesis and distribution of carotenoid in photosynthesis bacteria	11
Biosynthesis and distribution of carotenoid in algae and fungi	23
Regulation and control of carotenoids biosynthesis	27
Metabolism and distribution of carotenoids in aquatic animal	29
Biological function of carotenoid in aquatic animal	35
Applications of carotenoids for pigmentation in aquatic animal	41
Use of microorganism as a carotenoid source in aquatic animal feed	44
Photosynthetic bacteria and its applications in aquaculture	48
References	50
OBJECTIVES	60
CHAPTER II : BACTERIAL COLLECTION AND SCREENING FOR THE HIGHLY CAROTENOID PRODUCTION STRAIN	61
Introduction	61
Materials and methods	62
Results	66
Bacterial collection and screening for the highest carotenoids contents strain	66
Identification of the selected strain	66
16S rDNA sequencing and phylogenic analysis	72

Carotenoids analysis	80
----------------------	----

CONTENTS (Continued)

	page
Discussion	86
References	88

CHAPTER III : EFFECTS OF SYNTHETIC ASTAXANTHIN AND

PHOTOSYNTHETIC BACTERIAL CELLS ON GROWTH PERFORMANCE, PIGMENTATION AND IMMUNE RESPONSES IN BLACK TIGER SHRIMP (*PENAEUS MONODON*)

	91
Introduction	91
Materials and methods	92
Results	98
Pigmentation	98
Growth performances	98
Carotenoid analysis	103
Blood parameters	103
Histological changes	106
Discussion	109
References	112

CHAPTER IV : OPTIMIZATION FOR GROWTH AND TOTAL CAROTENOIDS

PRODUCTION OF *RHODOBACTOR* SP. TM11B FOR USE AS NUTRIENT SOURCE IN BLACK TIGER SHRIMP FEED

	115
Introduction	115
Materials and methods	116
Results	120
Effect of yeast extract/peptone (Y/P) supplementation	120
Effect of carbon source	121

Optimum concentration of monosodium glutamate for growth and total carotenoids production	124
Effects of metal ions supplement	126

CONTENTS (Continued)

	page
Effect of sodium chloride	132
Effects of pH	134
Carotenoids analysis	136
Discussion	137
References	140

CHAPTER V : THE EFFECTS OF *RHODOBACTER CAPSULATUS* TM11B ON GROWTH PERFORMANCE, PIGMENTATION, BLOOD PARAMETERS AND DISEASE RESISTANCE IN BLACK TIGER SHRIMP

	143
Introduction	143
Materials and methods	144
Results	151
Growth performances	151
Blood parameters	152
Carotenoid analysis	153
Stress tolerance	161
Disease resistance	162
Histological studies	163
Discussion	165
References	169

CHAPTER VI : Effect OF CAROTENOIDS ON HEALTH CONDITION, IMMUNE RESPONSE AND DISEASE RESISTANCE IN THE BLACK

TIGER SHRIMP	174
Introduction	174
Materials and methods	175
Results	182
Trial 1 Experiment in small shrimp	182
Trial 2 Experiment in juvenile shrimp	186
Discussion	190

CONTENTS (Continued)

	page
References	192
CHAPTER VII : EFFECT OF ALGRO NATURAL[®] ON GROWTH PERFORMANCE, HEALTH CONDITION, IMMUNE RESPONSE AND DISEASE RESISTANCE IN BLACK TIGER SHRIMP	195
Introduction	195
Materials and methods	196
Results	201
Trial 1 (small shrimp) : Growth performance	
Body color, Blood parameters and disease resistance	201
Trial 2 (juvenile shrimp) : Blood parameters	
Body colors and carotenoids analysis	203
Bacterial clearance ability	209
Discussion	210
References	213
CHAPTER IIX : CONCLUSION	216
APPENDIX	218
VITAE	249

LIST OF TABLES

	page
Table 1-1. Black tiger shrimp production (mt) from aquaculture in Thailand during 1995-2003	2
Table 1-2. Carotenogenesis pathways and their distribution within anaerobic photosynthetic bacteria	11
Table 2-1. Photosynthetic bacterial isolates with the capability for growth in either aerobic-dark and anaerobic light conditions used in the studies	67
Table 2-2. Total carotenoids (mg/g dry weight) of each isolates during 24-96 hrs. of the incubation periods under aerobic dark condition	69
Table 2-3. Total carotenoids (mg/g dry weight) of each isolates during 24-96 hrs. of the incubation periods under anaerobic light condition	71
Table 2-4. Physiological characteristics of cultures TM3, TN5, SV2 and TM11B	73
Table 3-1. Feed formulation of test diets	94
Table 3-2. Salmofan scores of the shrimp fed the experimental diets for 8 weeks	99
Table 3-3. Average body weight (g) of shrimp fed with each experimental diet during 8 weeks of feeding period	101
Table 3-4. Growth performance of the shrimp fed with each experimental diet for 8 weeks	102
Table 3-5. Total carotenoid, free astaxanthin, astaxanthin mono-ester and di-ester in shrimp fed with each experimental diet for 8 weeks	104
Table 3-6. Total hemocytes, phenoloxidase activity, superoxide dismutase activity and total antioxidant status of the shrimp fed with each experimental diet for 8 weeks	105
Table 3-7. Levels and frequency of histological changes (%) in black tiger	

shrimp fed with each experimental diet for 8 weeks 106

LIST OF TABLES (Continued)

	page
Table 5-1. Feed formulation of test diets.	150
Table 5-2. Average body weight (g), Weight gain (%), Survival (%) and FCR of the shrimp fed each experimental diet for 8 weeks periods.	152
Table 5-3. Blood parameter of the shrimp fed each experimental diet for 8 week.	153
Table 5-4. Total carotenoid and astaxanthin content in the shrimp fed each experimental diet for 8 weeks.	154
Table 6-1. Composition of the test diet for small shrimp in trial 1.	177
Table 6-2. Composition of the test diet for juvenile shrimp in trial 2.	178
Table 6-3. Analyzed of carotenoid compositions in each test diet for small shrimp in trial 1.	179
Table 6-4. Analyzed of carotenoid compositions in each test diet for juvenilel shrimp in trial 2. trial 2.	179
Table 6-5. Growth performance of small shrimp (avg. weigh 1-5 g) after feeding with each carotenoid pigments for 8 weeks.	183
Table 6-6. Salmo fan score, astaxanthin and total carotenoid in juvenile shrimp after fed each experimental diet for 6 weeks.	187
Table 6-7. Blood parameters and immune response of juvenile shrimp fed diet containing with carotenoid pigments for 6 weeks.	188
Table 7-1. Compositions of test diet.	200
Table 7-2. Average weigh (g) of small shrimp fed with experimental diet for 8 weeks.	202
Table 7-3. Growth performance of small shrimp after feeding with experimental diet for 8 weeks.	202
Table 7-4. Color score and visual apparent of the shrimp in each treatment after boiling for 5 min.	203
Table 7-5. Blood parameters of small shrimp fed experimental diet for 8 weeks	204

Table 7-6. Total hemocyte counts and phenoloxidase activity of juvenile shrimp fed experimental diet for 6 weeks	204
--	-----

LIST OF TABLES (Continued)

	page
Table 7-7. Salmo fan score and visual apparent of the juvenile shrimp fed experimental diet after boiling for 3 min.	205
Table 7-8. Total carotenoid and astaxanthin content in juvenile shrimp fed experimental diet for 6 weeks	209
Table 7-9. Total bacteria count in hemolymph after injected with suspension of <i>V. harveyi</i> for 3 hr.	210
Appendix Table 1. Cell mass (mg dry wt./ml) of PSB TM 11B in the medium contain each level of yeast extract and peptone	219
Appendix Table 2. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM 11B in the medium contain each level of yeast extract and peptone.	220
Appendix Table 3. Cell mass (mg dry wt./ml) of PSB TM11B in the medium contained each carbon source (4g/l glutamic acid as nitrogen source in the basal medium)	221
Appendix Table 4. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM11B in the medium contained each carbon source (4g/l glutamic acid as nitrogen source in the basal medium).	222
Appendix Table 5. Cell mass (mg dry wt./ml) of PSB TM11B in the medium contained glutamic acid and monosodium glutamate as C and N source.	223
Appendix Table 6. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM11B in the medium contained glutamic acid and monosodium glutamate as C and N source	224
Appendix Table 7. Cell mass (mg dry wt./ml) of PSB TM11B in the medium	

contained each concentration of mono sodium glutamate. 225

LIST OF TABLES (Continued)

	page
Appendix Table 8. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM 11B in the medium contained each concentration of mono sodium glutamate.	226
Appendix Table 9. Cell mass (mg dry wt./ml) of PSB TM11B in the medium contained each concentration of metal ions.	227
Appendix Table 10. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM 11B in the medium contained each concentration of metal ions.	228
Appendix Table 11. Cell mass (mg dry wt./ml) of PSB TM11B in the medium contained each concentration of cobalt chloride.	229
Appendix Table 12. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM 11B in the medium contained each concentration of cobalt chloride.	230
Appendix Table 13. Cell mass (mg dry wt./ml) of PSB TM 11B in the medium contained each concentration of ferric chloride.	231
Appendix Table 14. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM 11B in the medium contained each concentration of ferric chloride.	232
Appendix Table 15. Cell mass (mg dry wt./ml) of PSB TM 11B in the medium contained each concentration of ferric citrate.	233
Appendix Table 16. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM 11B in the medium contained each concentration of ferric citrate.	234
Appendix Table 17. Cell mass (mg dry wt./ml) of PSB TM 11B in the medium contained each concentration of sodium chloride.	235
Appendix Table 18. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM 11B in the medium contained each concentration of sodium chloride.	236
Appendix Table 19. Cell mass (mg dry wt./ml) of PSB TM 11B in the difference pH of the culture medium.	237

Appendix Table 20. Total carotenoids contents ($\mu\text{g/g}$ dry wt.) of PSB TM 11B in the the difference pH of the culture medium.	238
---	-----

LIST OF FIGURES

	page
Figure 1-1. Isoprene unit	5
Figure 1-2. Lycopene	6
Figure 1-3. β -carotene	6
Figure 1-4. Formation of geranylgeranyl pyrophosphate (GGPP) by the basic isoprenoid biosynthesis pathway from acetyl-CoA	8
Figure 1-5. General isoprenoid biosynthesis pathway and the mechanism for the formation of phytoene	9
Figure 1-6. Desaturation of phytoene to lycopene by phytoene desaturase	12
Figure 1-7. The predicted pathway for the biosynthesis of spirilloxanthin	14
Figure 1-8. The pathway for the biosynthesis of spheroidene	15
Figure 1-9. The predicted pathway for the biosynthesis of cross-conjugated carotenals and their structure	17
Figure 1-10. The predicted pathway for the biosynthesis of isorenieratene and Chlorobactene	19
Figure 1-11. Bioconversion pathways of carotenoids in the penaeid shrimp and proposed conversion pathway to retinoids	37
Figure 2-1. The first 500bp DNA sequence of the 16S rDNA from strain TM3	74
Figure 2-2. The first 500bp DNA sequence of the 16S rDNA from strain TM11 B	74
Figure 2-3. The first 500bp DNA sequence of the 16S rDNA from strain SV2	75

Figure 2-4. The first 500bp DNA sequence of the 16S rDNA from strain TN5	75
Figure 2-5. The phylogenic tree of strain TM 3	76
Figure 2-6. The phylogenic tree of strain TM 11 B	77
Figure 2-7. The phylogenic tree of strain SV 2	78

LIST OF FIGURES (Continued)

	page
Figure 2-8. The phylogenic tree of strain TN 5	79
Figure 2-9. Thin layer chromatogram of pigments extracted from 4 strains of photosynthetic bacteria	80
Figure 2-10. VIS spectrum (upper) and mass spectrum (lower) of band A which corresponds to neurosporene	81
Figure 2-11. VIS spectrum (upper) and mass spectrum (lower) of band B which corresponds to spheroidene	82
Figure 2-12. VIS spectrum (upper) and mass spectrum (lower) of band C which corresponds to spheroidenone	83
Figure 2-13. VIS spectrum (upper) and mass spectrum (lower) of band D which corresponds to demethylspheroidene	84
Figure 2-14. VIS spectrum (upper) and mass spectrum (lower) of band E which corresponds to hydroxyspheroidene	85
Figure 3-1. Body color of shrimp fed with each experimental diet for 8 weeks	108
Figure 3-2. Color of boiled shrimp fed each experimental diet	100
Figure 3-3. Atrophy in tubular epithelium of hepatopancreas, cell necrosis and degeneration of tubules with the infiltration of haemocytes during hepatopancreatic tubues, epithelium cells of antennal gland develop signs of atrophy and degeneration of intestinal epithelium in shrimp fed with <i>Rhodobactes sphaeroides</i> SV2	108
Figure 4-1. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each level of Y/P after cultivation for 96 hr.	120

Figure 4-2. Cell mass (mg/ml) (a) and total carotenoids (b) of PSB TM 11B in medium contain each level of Y/P after cultivation for 120 hr.	121
--	-----

LIST OF FIGURES (Continued)

	page
Figure 4-3. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each carbon source after cultivation for 96 hr.	122
Figure 4-4. Cell mass (mg/ml) (a) and total carotenoids (b) of PSB TM 11B in medium contain each carbon source after cultivation for 120 hr.	122
Figure 4-5. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain glutamic acid and mono sodium glutamate after cultivation for 96 hr.	123
Figure 4-6. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain glutamic acid and mono sodium glutamate after cultivation for 120 hr.	124
Figure 4-7. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each level of mono sodium glutamate after cultivation for 96 hr.	125
Figure 4-8. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each level of mono sodium glutamate after cultivation for 120 hr.	125
Figure 4-9. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain ferrous sulphate and magnesium sulphate after cultivation for 96 hr.	126
Figure 4-10. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor</i> <i>capsulatus</i> TM11B in medium contain ferrous sulphate and magnesium sulphate after cultivation for 120 hr.	127

Figure 4-11. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain ferrous sulphate and magnesium sulphate after cultivation for 96 hr.	128
--	-----

LIST OF FIGURES (Continued)

	page
Figure 4-12. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain ferrous sulphate and magnesium sulphate after cultivation for 96 hr.	128
Figure 4-13. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each concentration of ferric chloride after cultivation for 96 hr.	129
Figure 4-14. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each concentration of ferric chloride after cultivation for 120 hr.	130
Figure 4-15. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each concentration of ferric citrate after cultivation for 96 hr.	131
Figure 4-16. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each concentration of ferric citrate after cultivation for 120 hr.	131
Figure 4-17. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each concentration of sodium chloride after cultivation for 96 hr.	133
Figure 4-18. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in medium contain each concentration of sodium chloride after cultivation for 120 hr.	133
Figure 4-19. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobactor capsulatus</i> TM11B in the difference pH of culture medium after	

cultivation for 96 hr	134
Figure 4-20. Cell mass (mg/ml) (a) and total carotenoids (b) of <i>Rhodobacter capsulatus</i> TM11B in the difference pH of culture medium after cultivation for 120 hr.	135

LIST OF FIGURES (Continued)

	page
Figure 4-21. Thin layer chromatogram of carotenoids pigment from <i>Rhodobacter capsulatus</i> TM11B in the G5 (right) or optimized medium used in this study (left).	136
Figure 5-1. Body color of shrimp fed each test diet for 8 weeks.	155
Figure 5-2. Body color of shrimp fed each test diet after boiling for 3 min.	155
Figure 5-3. Thin layer chromatogram of carotenoids extracted from the shrimp fed test diet for 8 weeks	156
Figure 5-4. VIS spectrum of the pigment A (β -carotene) extracted from the shrimp fed diet 1-6 for 8 weeks compared to the standard UV/VIS spectrum	157
Figure 5-5. VIS spectrum of the pigment B (diester astaxanthin) extracted from the shrimp fed diet 1-6 for 8 weeks compared to the standard UV/VIS spectrum	158
Figure 5-6. VIS spectrum of the pigment C (mono-ester astaxanthin) extracted from the shrimp fed diet 1-6 for 8 weeks.	159
Figure 5-7. VIS spectrum of the pigment D (free- astaxanthin) extracted from the shrimp fed diet 1-6 for 8 weeks	160
Figure 5-8. Survival rates (%) of the shrimp in each treatment during 7-days of stress period.	161
Figure 5-9. Survival rates (%) of the shrimp in each treatment after challenged with <i>Vibrio harveyi</i> for 14 days.	162
Figure 5-10. (a) Hepatopancreas of the shrimp fed diet supplemented	

with 5% <i>Rhodobacter</i> sp. TM11B (b) Antennal gland of the shrimp fed diet supplemented with 5% <i>Rhodobacter</i> sp. TM11B	
(c) Intestinal epithelium of the shrimp fed diet supplemented with 5% <i>Rhodobacter</i> sp. TM11B	164

LIST OF FIGURES (Continued)

	page
Figure 6-1. Survival rates of small shrimp fed diet containing with each carotenoid pigments after challenged with WSSV at a concentration of 1 : 10 ⁷ for 13 days.	184
Figure 6-2. Survival rates of small shrimp fed diet containing with each carotenoid pigments during salinity stress for 13 days.	185
Figure 6-3. Survival rates of juvenile shrimp fed diet containing with each carotenoid pigments after challenged with WSSV at a concentration of 1 : 10 ⁷ for 10 days.	189
Figure 7-1. Survival rates (%) of the shrimp in each treatment after challenged with 1:10 ⁶ WSSV for 15 days.	206
Figure 7-2. Survival rates (%) of the shrimp in each treatment during 9-days of stress period	207
Figure 7-3. Body color of black tiger shrimp fed each test diet for 8 weeks.	208