

REFERENCES

- A.O.A.C. 1999. *Official Methods of Analysis*. 14th ed. Washington, D. C: Association of Agricultural Chemists.
- Abrahamson, M. Mason, R. and Hansson, H. 1991. Human cystatin C: role of the N-terminal segment in the inhibition of human cysteine proteinases and in its inactivation by leukocyte elastase. *J. Biochem.* 273: 621-626.
- Adam, A., Albert, A., Calay, G., Closset, J., Damas, J. and Franchimont, P. 1985. Human kininogen of low molecular mass: quantification by radioimmunoassay and determination of reference value. *Clin. Chem.* 31: 423-426.
- Akazawa, H., Miyauchi, Y., Sakurada, K., Wasson, D. H. and Reppond, K. D. 1993. Evaluation of protease inhibitors in Pacific whiting surimi. *J. Aquat. Food Prod. Technol.* 2 (3): 79-95.
- Akpinar, O. and An, H., 2005. Purification and determination of inhibitory activity of recombinant soyacystatin for surimi application. *Mol. Nutr. Food Res.* 49: 247-255.
- Alvarez, C., Couso, I., Solas, M. and Tejada, M. 1997. Waxy corn starch affecting texture and ultrastructure of sardine surimi gels. *Z. Lebensm Unters Forsch A.* 204: 121-128.
- Alvarez, C., Couso, I. and Tejada, M. 1999. Thermal gel degradation (modori) in sardine surimi gels. *J. Food Sci.* 64: 633-637.
- An, H., Peters, M. Y., Seymour, T. A. and Morrissey, M. T. 1995. Isolation and activation of cathepsin-L inhibitor complex from Pacific whiting (*Merluccius productus*). *J. Agric. Food Chem.* 43: 327-330.
- An, H., Peters, M. Y., and Seymour, T. A. 1996. Roles of endogenous enzymes in surimi gelation. *Trend Food Sci. Technol.* 7: 321-327.
- An, H., Seymour, T. A., Wu, J and Morrissey, M. T. 1994a. Assay system and characterization of Pacific whiting (*Merluccius productus*) protease. *J. Food Sci.* 59: 227-281.
- An, H., Weerasinghe, V., Seymour, T. A. and Morrissey, M. T. 1994b. Cathepsin degradation of Pacific whiting surimi proteins. *J. Food Sci.* 59: 1013-1017, 1033.

- Anastasi, A., Brown, M. A., Kembhavi, A. A., Nicklin, M. J. H., Sayers, C. A., Sunter, D. C. and Barrett, A. J. 1983. Cystatin, a protein inhibitor of cysteine proteinases. *Biochem. J.* 211, 129–138.
- Aoki, T. and Ueno, R. 1997. Involvement of cathepsin B and L in the post-mortem autolysis of mackerel muscle. *Food Res. Int.* 30: 585–591
- Aranishi, F., Ogata, H., Hara, K., Osatomi, K. and Ishihara, T. 1998. Susceptibility of opioid peptides and myofibrillar proteins to carp cathepsin L. *J. Agric. Food Chem.* 46: 388–392.
- Asher, D. M. 1999. Bovine sera used in the manufacture of biologicals: current concern and policies of the U.S Food and Drug Administration regarding the Transmissible Spongiform Encephalopathies. *Develop. Biol. Stand.* 99: 41–44.
- Ashie, I. N. A. and Simpson, B. K. 1997. Proteolysis in food myosystems—a review. *J. Food Biochem.* 21: 91–123.
- Ashie, I. N. A. and Lanier, T. C. 2000. Transglutaminase in seafood processing. *In* *Seafood Enzymes: Utilization and Influence on Post-Harvest Seafood Quality.* (Haard, N. F., and Simpson, B. K. eds.). p. 147–166). Marcel Dekker, Inc. New York.
- Barrett, A. J. 1995. Classification of peptidases: proteolytic enzymes: serine and cysteine peptidases. *Methods Enzymol.* 244: 1–15.
- Barrett, A. J., Fritz, H., Grubb, A., Isemura, S., Jarvinen, M., Katunuma, N., Machleidt, W., Muller-Esterl, W., Sasaki, M. and Turk, V. 1986. Nomenclature and classification of the proteins homologous with the cysteine–protease inhibitor chicken cystatin. *Biochem. J.* 236: 321.
- Barrett, A. J. and McDonald, J. K. 1980. *Mammalian proteinases: A glossary and bibliography.* Vol. 1. Academic Press. London.
- Barrett, A. J., Rawlings, N. D., Davies, M. E., Machleidt, W., Salvesen, G. and Turk, V. 1986. Cysteine proteinase inhibitors of the cystatin superfamily. *In* *Proteinase Inhibitors.* (Barrett, A. J. and Salvesen, G. eds.). p. 515–569. Elsevier Science Publishers. Amsterdam.
- Barrett, A. J. and Starkey, P. M. 1973. The interaction of α_2 -macroglobulin with proteinases: characteristics and specificity of the reaction, and a hypothesis concerning its molecular mechanism. *Biochem J.* 133: 709–724.

- Benjakul, S., Chantarasuwan, C. and Visessanguan, W. 2003b. Effect of medium temperature setting on gelling characteristics of surimi from some tropical fish. *Food Chem.* 84: 567–574.
- Benjakul, S., Leelapongwattana, K. and Visessanguan, W. 2003a. Purification and characterization of heat-stable alkaline proteinase from bigeye snapper (*Priacanthus macracanthus*) muscle. *Comp. Biochem. Physiol.* 134B: 579–591.
- Benjakul, S., Seymour, T. A., Morrissey, M. T. and An, H. 1996. Proteinase in Pacific whiting surimi wash water: identification and characterization. *J. Food Sci.* 61: 1165–1170.
- Benjakul, S., Seymour, T. A., Morrissey, M. T. and An, H. 1997a. Physicochemical changes in Pacific whiting muscle proteins during iced storage. *J. Food Sci.* 62: 729–733.
- Benjakul, S., Seymour, T. A., Morrissey, M. T. and An, H. 1997b. Recovery of proteinase from Pacific whiting surimi wash water. *J. Food Biochem.* 21: 431–443.
- Benjakul, S., Seymour, T. A., Morrissey, M. T. and An, H. 1998. Characterization of proteinase from Pacific whiting surimi wash water. *J. Food Biochem.* 22: 1–16.
- Benjakul, S. and Visessanguan, W. 2000. Pig plasma protein: potential use as proteinase inhibitor for surimi manufacture; inhibitory activity and the active components. *J. Sci. Food Agric.* 80: 1351–1356.
- Benjakul, S. and Visessanguan, W. 2003. Transglutaminase-mediated setting in bigeye snapper surimi. *Food Res. Int.* 36: 253–266.
- Benjakul, S., Visessanguan, W. and Chantarasuwan, C. 2004a. Cross-linking activity of sarcoplasmic fraction from bigeye snapper (*Priacanthus tayenus*) muscle. *Lebensm.-Wiss. u.-Technol.* 37: 79–85.
- Benjakul, S., Visessanguan, W., Ishizaki, S. and Tanaka, M. 2001d. Differences in gelation characteristics of natural actomyosin from two species of bigeye snapper, *Priacanthus tayenus* and *Priacanthus macracanthus*. *J. Food Sci.* 66: 1311–1318.
- Benjakul, S., Visessanguan, W. and Srivilai, C. 2001a. Porcine plasma protein as gel enhancer in bigeye snapper (*Priacanthus tayenus*) surimi. *J. Food Biochem.* 25: 285–305.

- Benjakul, S., Visessanguan, W. and Srivilai, C. 2001b. Gel properties of bigeye snapper (*Priacanthus tayenus*) surimi as affected by setting and porcine plasma proteins. *J. Food Quality*. 24: 453–472.
- Benjakul, S., Visessanguan, W. and Srivilai, C. 2001c. Porcine plasma protein as proteinase inhibitor in bigeye snapper (*Priacanthus tayenus*) muscle and surimi. *J. Sci. Food Agric*. 81: 1039–1046.
- Benjakul, S., Visessanguan, W. and Turksuban, J. 2003c. Changes in physico-chemical properties and gel-forming ability of lizardfish (*Saurida tumbil*) during post-mortem storage in ice. *Food Chem*. 80: 535–544.
- Benjakul, S., Visessanguan, W. and Turksuban, J. 2003d. Heat-activated proteolysis in lizardfish (*Saurida tumbil*) muscle. *Food Res. Int*. 36: 1021–1028.
- Benjakul, S., Visessanguan, W., Tueksuban, J. and Tanaka, M. 2004b. Effect of some protein additives on proteolysis and gel forming ability of lizardfish (*Saurida tumbil*). *Food Hydrocolloid*. 18: 395–401.
- Boye, S. W. and Lanier, T. C. 1988. Effects of heat-stable alkaline proteinase activity of Atlantic menhaden (*Brevoorti tyrannus*) on surimi gels. *J. Food Sci*. 53: 1340–1342, 1398.
- Bradley, R. 1999. BSE transmission studies with particular reference to blood. *Develop. Biol. Stand*. 99: 35–40.
- Brzin, J., Popovic, T. and Turk, V. 1984. Human cystatin, a new protein inhibitor of cysteine proteinases. *Biochem. Biophys. Res. Comm*. 118: 103–109.
- Burgarella, J. C., Lanier, T. C., Hamann, D. D. and Wu, M. C. 1985. Gel strength development during heating of surimi in combination with egg white or whey protein concentrate. *J. Food Sci*. 50: 1595–1597.
- Burnouf, T. 1995. Chromatography in plasma fractionation: benefits and future trends. *J. Chromatogra B*. 664: 3–15.
- Caldironi, H. A. and Ockerman, H. W. 1982. Incorporation of blood proteins into sausage. *J. Food Sci*. 47: 405–408.
- Cao, M. J., Hara, K., Osatomi, K., Tachibana, K., Lzumi, T. and Ishihara, T. 1999. Myofibril-bound serine proteinase (MBP) and its degradation of myofibrillar proteins. *J. Food Sci*. 64: 644–647.
- Cao, M. J., Osatomi, K., Hara, K. and Ishihara, T. 2000. Identification of a myofibril-bound serine proteinase (MBSP) in the skeletal muscle of lizardfish (*Saurida*

- wanieso*) which specifically cleaves the arginine site. *Comp. Biochem. Physiol.* 125B: 255–264.
- Carlson, T. H. 1996. Composition of plasma. *In Principles of Transfusion Medicine*. 2nd ed. (Rossi, E. C., Simon, T. L., Moss, G. S. and Gould, S. A., eds.). p. 383–391. Williams & Wilkins. Baltimore.
- Careche, M., Alvarez, C. and Tejada, M. 1995. Suwari and kamaboko gels: effect of heat treatment on solubility of networks. *J. Agric. Food Chem.* 43: 1002–1010.
- Carvajal, P. A. 2002. Diffusion of active proteins into fish meat to minimize proteolytic degradation. Ph. D. Dissertation. North Carolina State University.
- Chang-Lee, M. V., Lampila, L. E. and Crawford, D. L. 1990. Yield and composition of surimi from Pacific whiting (*Merluccius productus*) and the effect of various protein additives on gel strength. *J. Food Sci.* 55: 83–86.
- Chang-Lee, M. V., Pacheco-Aguilar, D. L., Crawford, L. and Lampila, L. E. 1989. Proteolytic activity of surimi from Pacific whiting (*Merluccius productus*) and heat-set gel texture. *J. Food Sci.* 54: 1116–1119, 1124.
- Ching, H. L. and Leinot, A. 1993. Surimi composition and technology: present status and nutritional considerations. *Int. J. Food Sci. Nutr.* 44, 555–563.
- Choi, Y. J., Cho, Y. J. and Lanier, T. C. 1999. Purification and characterization of proteinase from Atlantic menhaden muscle. *J. Food Sci.* 64: 772–775.
- Choi, Y. J., Kang, I. S. and Lanier, T. C. 2004. Proteolytic enzymes and control in surimi. *In Surimi and Surimi Seafood*. 2nd ed. (Park, J. W., ed.). p. 227–278. Taylor & Francis. New York.
- Chung, K. H. and Lee, C. M. 1991. Water binding and ingredient dispersion pattern effects on surimi gel texture. *J. Food Sci.* 56: 1263–1266.
- Cohen, I., Young-Bandala, L., Blankenberg, T. A., Siefring, G. E. and Bruner-Lorand, J. 1979. Fibrinolytic-catalyzed cross-linking of myosin from platelet and skeletal muscle. *Arch. Biochem. Biophys.* 192: 100–111.
- Cofrades, S., Guerra, M. A., Carballo, J., Fernandez-Martin, F. and Jimenez-Colmenero, F. 2000. Plasma protein and soy fiber content effect on bologna sausage properties as influenced by fat level. *J. Food Sci.* 65: 281–287.
- Cohn, E. J., Strong, L. E., Hughes, W. L., Mulford, D. J., Ashworth, J. N., Melin, M. and Taylor, H. L. 1946. Preparation and properties of serum and plasma proteins.

- IV. A system for the separation into fraction of the protein and lipoprotein components of biological tissues and fluids. *J. Am. Chem. Soc.* 68: 459–475.
- Cooke, R. D. and Holbrook, J. J. 1974. Calcium and the assays of human plasma clotting factor XIII. *Biochem. J.* 141: 71–78.
- Credo, R. B., Curtis, C. G. and Lorand, L. 1978. Ca²⁺-related regulatory function of fibrinogen. *In* Proceeding of the National Academy of Sciences of the United States of American. 75: 4234–4237.
- Christeller, J. T., Farley, P. C., Ramsay, R. J., Sullivan, P. A. and Laing, W. A. 1998. Purification, characterization and cloning of an aspartic proteinase inhibitor from squash phloem exudate. *Eur. J. Biochem.* 254: 160–167.
- Chung, K. H. and Lee, C. M. 1991. Water binding and ingredient dispersion pattern effects on surimi texture. *J. Food Sci.* 56: 1263–1266.
- Curling, J. M. 1980. *Methods of plasma protein fractionation.* Academic Press. New York.
- Davey, C. L. and Gilbert, K. V. 1969. Studies in meat tenderness: changes in the fine structure of meat during aging. *J. Food Sci.* 34: 69–72.
- De Backer-Royer, C., Traore, F. and Meunier, J. C. 1992. Purification and properties of factor XIII from human placenta. *Int. J. Biochem.* 24: 91–97.
- Delaney, R. A. M. 1975. The nutritive value of porcine blood plasma concentrates prepared by ultra filtration and spray drying. *J. Sci. Food Agric.* 26: 303–310.
- Department of Livestock Development. 1996. *Data of livestock economic for the year 1996.* Bangkok: Thailand.
- Department of Livestock Development. 2004. *Statistics of import/export for chicken meat and product.* Information and Statistics Group, Information Technology Center, Ministry of Agricultural and Cooperatives, Bangkok: Thailand.
- Dill, K. A. and Shortle, D. 1991. Denatured states of proteins. *Ann. Rev. Biochem.* 60: 795–825.
- Duarte, R. T., Carvalho Simoes, M. C. and Sgarbieri, V. C. 1999. Bovine blood components: fractionation, composition, and nutritive value. *J. Agric. Food Chem.* 47: 231–236.
- Egelandsdal, B., Fretheim, K. and Samejima, K. 1986. Dynamic rheological measurements on heat-induced myosin gels: effect of ionic strength, protein concentration and addition of adenosine triphosphate or pyrophosphate. *J. Sci. Food Agric.* 37: 915–926.

- Fernando, T. 1992. Blood meal, meat and bone meal and tallow. *In* Inedible Meat By-products Advance in Meat Research (Pearson, A. M. and Dutson, T. R., eds.). p. 81-112. Elsevier Applied Science. London.
- Fishman, W. 1960. Plasma enzymes. *In* The Plasma Proteins. vol. II. (Putnum, F., ed.). p. 59-103. Academic Press. New York.
- Foegeding, E. A., Dayton, W. R. and Allen, C. E. 1986. Interaction of myosin-albumin and myosin-fibrinogen to form protein gels. *J. Food Sci.* 51: 109-112.
- Folk, J. E. 1980. Transglutaminase. *Ann. Rev. Biochem.* 49: 517-531.
- Gaertner, H. F. and Puigserver, A. J. 1992. Increased activity and stability of poly(ethylene glycol)-modified trypsin. *Enz. Microbiol. Technol.* 14: 150-155.
- Garcia-Carreno, F. L. 1996. Proteinase inhibitors. *Trend Food Sci. Technol.* 7: 197-204.
- Garcia-Carreno, F. L., Dimes, L. E. and Haard, N. F. 1993. Substrate-gel electrophoresis for composition and molecular weight of proteinases or proteinaceous proteinase inhibitors. *Anal. Biochem.* 214: 65-69.
- Garcia-Carreno, F. L. and Hernandez-Cortes, P. 2000. Use of protease inhibitors in seafood products. *In* Seafood Enzymes. (Haard, N. F. and Simpson, B. K., eds.). p. 531-547. Marcel Dekker, Inc. New York.
- Gautier, J., Bird, D. M. and Ng Kwai Hang, K. F. 1983. Plasma protein patterns in captive American kestrels. *Comp. Biochem. Physiol.* 75: 163-166.
- Gomez-Guillen, M. C., Borderias, A. J. and Montero, P. 1996. Rheological properties of gels made from high and low quality sardine (*Sardina pilchardus*) mince with added nonmuscle proteins. *J. Agric. Food Chem.* 44: 746-750.
- Gomez-Guillen, M. C., Borderias, A. J. and Montero, P. 1997. Chemical interactions of non-muscle proteins in the network of sardine (*Sardina pilchardus*) muscle gels. *Lebensm.-Wiss. u.-Technol.* 29: 602-608.
- Gomez-Guillen, M. C., Hurtado, J. L and Montero, P. 2002. Autolysis and protease inhibition effects on dynamic viscoelastic properties during thermal gelation of squid muscle. *J. Food Sci.* 67: 2491-2496.
- Gomez-Guillen, M. C. and Montero, P. 1996. Addition of hydrocolloids and non-muscle proteins to sardine (*Sardina pilchardus*) mince gels: effect of salt concentration. *Food Chem.* 56: 421-427.

- Gorman, J. J. and Folk, J. E. 1980. Structural features of glutamine substrates for human plasma factor XIIIa (activated blood coagulation factor XIII). *J. Biol. Chem.* 255: 419-427.
- Guenneugues, P. and Morrissey, M. T. 2004. Surimi resources. *In* Surimi and Surimi Seafood. 2nd ed. (Park, J. W., ed.). p. 3-32. Taylor & Francis Group. New York.
- Hall, G. M. and Ahmad N. H. 1997. Surimi and fish-mince products. *In* Fish Processing Technology. 2nd ed. (Hall, G. M., ed.). p. 74-92. Blackie academic & Professional. London.
- Hamann, D. D. (1988). Rheology as a means of evaluating muscle functionality of processed foods. *Food Technol.* 42(6): 66-71.
- Hamann, D. D., Amato, P. M., Wu, M. C. and Foegeding, E. A. 1990. Inhibition of modori (gel weakening) in surimi by plasma hydrolysate and egg white. *J. Food Sci.* 55: 665-669, 795.
- Hamann, D. D. and MacDonald, G. A. 1992. Rheology and texture properties of surimi and surimi-based foods. *In* Surimi Technology. (Lanier, T. C. and Lee, C. M., eds.). p. 429-500. Marcel Dekker. New York.
- Hao, Y. L., Ingham, K. C. and Wickerhauser, M. 1980. Fractional precipitation of proteins with polyethylene glycol. *In* Methods of Plasma Protein Fractionation. (Curling, J. M., ed.). p. 57-76. Academic Press. London.
- Harpel, P. C. and Brower, M. S. 1983. α_2 -macroglobulin: an introduction. *In* Chemistry and Biology of α_2 -macroglobulin. (Boland, B., Hitchcock, J. and Scholnick, R.E. eds.). p. 1-9. The N.Y. Acad. Sci. New York.
- Hasko, F. and Vassilyeva, R. 1981. Fractionation of plasma protein with PEG. *Haematologic (Budap).* 14: 199-206.
- Hayashi, I., Kato, H., Iwanaga, S. and Oh-ishi, S. 1985. Rat plasma high-molecular weight kininogen. A simple method for purification and its characterization. *J. Biol. Chem.* 260: 6115-6123.
- Heu, M. S., Kim, H. R., Cho, D. M., Godber, J. S. and Pyeun, J. H. 1997. Purification and characterization of cathepsin L-like enzyme from muscle of anchovy (*Engraulis japonica*). *Comp. Biochem. Physiol.* 118B: 523-529.
- Howell, N. K. and Lawrie, R. A. 1983. Functional aspects of blood plasma proteins: I Separation and characterization. *J. Food Technol.* 18: 747-762.

- Howell, N. K. and Lawrie, R. A. 1984. Functional aspects of blood plasma proteins: II Gelling properties. *J. Food Technol.* 19: 289–295.
- Hultin, H. O. and Kelleher, S. D. 2000. Surimi processing from dark muscle fish. *In* Surimi and Surimi Seafood. (Park, J. W., ed.). p. 59–77. Marcel Dekker. New York.
- Imai, C., Tsukamasa, Y., Sugiyama, M., Minegishi, Y. and Shimizu, Y. 1996. The effect of setting temperature on the relationship between ϵ -(γ -glutamyl) lysine cross-link content and breaking strength in salt-ground meat of sardine and Alaska pollack. *Nippon Suisan Gakkaishi.* 62: 104–111.
- Iso, N., Mizuno, H., Saito, T., Lin, C. Y., Fujita, T. and Nagaisha, E. 1985. The effect of additives (egg white and soy protein) on the rheological properties of kamaboko. *Nippon Suisan Gakkaishi.* 51: 485–488.
- Jiang, S. T. 2000. Enzymes and their effects on seafood texture. *In* Seafood Enzymes. (Haard N. F. and Simpson, B. K., eds.). p. 411–450. Marcel Dekker, Inc. New York.
- Jiang, L., He, L. and Fountoulakis, M. 2004. Comparison of protein precipitation methods for sample preparation prior to proteomic analysis. *J. Chromatography A.* 1023: 317–320.
- Jiang, S. T. and Lee, J. J. 1992. Purification, characterization and utilization of pig plasma factor XIIIa. *J. Agric. Food Chem.* 40: 1101–1107.
- Jiang, S. T., Lee, J. J. and Chen, H. C. 1996. Proteolysis of actomyosin by cathepsins B, L, L-like, and X from mackerel (*Scomber australasicus*). *J. Agric. Food Chem.* 44: 769–773.
- Kahn, D. R. and Cohen, I. 1981. Factor XIIIa-catalyzed coupling of structural proteins. *Biochim. Biophys. Acta.* 668: 490–494.
- Kang, I. S. and Lanier, T. C. 1997. Bovine plasma protein functions in surimi gelation compared with cysteine protease inhibitors. *J. Food Sci.* 64: 842–846.
- Kang, I. S. and Lanier, T. C. 1999. Bovine plasma protein functions in surimi gelation compared with cysteine protease inhibitors. *J. Food Sci.* 64: 842–846.
- Kang, I. S. and Lanier, T.C. 2000. Heat-induced softening of surimi gels by proteinases. *In* Surimi and Surimi Seafood. (Park, J. W., ed.). p. 445–474. Marcel Dekker, Inc. New York.

- Karlsrud, T. S., Bu, L., Aasen, A. O. and Johansen, H. T. 1996. Quantification of kininogen in plasma. A functional method based on the cysteine proteinase inhibitory activity. *Thromb. Res.* 82: 265-273.
- Kato, H., Nagasawa, S. and Iwanaga, S. 1981. HMW and LMW kininogens. *Methods Enzymol.* 80: 172-198.
- Kennedy, A. R. 1998. Chemopreventive agents: protease inhibitors. *Pharmacol. Ther.* 78: 167-209.
- Kent, R. S. and Drohan, W. N. 2001. Methods for the selective separation of organic components from biological fluids. U. S. Patent 6,193,891.
- Kim, S. H., Carpenter, J. A., Lanier, T. C. and Wicker, L. 1993. Polymerization of beef actomyosin induced by transglutaminase. *J. Food Sci.* 58: 473-474, 491.
- Kimura, M., Sueyoshi, T., Takada, K., Tanaka, K., Morita, T. and Iwanaga, S. 1987. Isolation and characterization of ornitho-kininogen. *J. Biochem.* 168: 493-501.
- Kimura, I., Sugimoto, M., Toyoda, K., Seki, N., Arai, K. and Fujita, T. 1991. A study on the cross-linking reaction of myosin in kamaboko (suwari) gels. *Nippon Suisan Gakkaishi.* 57: 1386-1396.
- Kinoshita, M., Toyohara, H. and Shimizu, Y. 1990. Purification and properties of a novel latent proteinase showing myosin heavy chain-degrading activity from threadfin bream muscle. *J. Food Biochem.* 107: 587-591.
- Kishi, H., Nozawa, H. and Seki, N. 1991. Reactivity of muscle transglutaminase on carp myofibrils and myosin B. *Nippon Suisan Gakkaishi.* 57: 1203-1210.
- Komiya, M., Kato, H. and Suzuki, T. 1974. Bovine plasma kininogens. Further purification of high molecular weight kininogen and its physicochemical properties. *J. Biochem.* 76: 811-822.
- Koohmaraie, M. 1989. The role of endogenous protease in meat tenderness. *Proc. Recip. Meat Conf.* 41: 89-100.
- Koohmaraie, M. 1994. Muscle proteinase and meat aging. *Meat Sci.* 36: 93-104.
- Kos, J., Dolinar, M. and Turk, V. 1992. Isolation and characterization of chicken L- and H-kininogens and their interaction with chicken cysteine proteinases and papain. *Agents Actions. Supple.* 38: 331-339.
- Kumazawa, Y., Numazawa, T., Seguro, K. and Motoki, M. 1995. Suppression of surimi setting by transglutaminase inhibitors. *J. Food Sci.* 60: 715-717.

- Kubota, M., Kinoshita, M., Kubota, S., Yamashita, M., Toyohara, H. and Sakaguchi, M. 2001. Possible implication of metalloproteinases in post-mortem tenderization of fish muscle. *Fish. Sci.* 67: 965–968.
- Kurth, L. 1983. Grosslinking of myosin and casein by the enzyme transglutaminase. *Food Technol. Aus.* 35: 420–424.
- Kuryl, J. and Gasparska, J. 1985. The differences in plasma protein pattern between laying and non-laying chickens, quails and geese. *Comp. Biochem. Physiol.* 80: 309–313.
- Ladrat, C., Chaplet, M., Verrez-Bagnis, V., Noel, J. and Fleurence, J. 2000. Neutral calcium-activated proteases from European sea bass (*Dicentrarchus labrax L*) muscle: polymorphism and biochemistry studies. *Comp. Biochem. Physiol.* 125B: 83–95.
- Ladrat, C., Verrez-Bagnis, V. and Fleurence, J. 2003. *In vitro* proteolysis of myofibrillar and sarcoplasmic proteins of white muscle of sea bass (*Dicentrarchus latex L*): effects of cathepsin B, D and L. *Food Chem.* 81: 517–525.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage. *Nature.* 227: 680–685.
- Lamb-Sutton, Y. 1995. The possible role of alpha-2-macroglobulin in the inhibition of fish muscle heat-stable proteases. M. S. Thesis. North Carolina State University.
- Lanier, T. C. 1986. Functional properties of surimi. *Food Technol.* 40(3): 107–114.
- Lanier, T. C. 1990. Interaction of muscle and non-muscle proteins affecting heat-set gel rheology. *In* *Macromolecular Interactions and Food Colloid Stability*. (Parris, N., Barford, R. A., eds.). p. 268–284. ACS Symposium Series. New York.
- Lanier, T. C. 1991. Interactions of muscle and non-muscle protein affecting heat set gel rheology. *In* *Interactions of Food Protein*. (Parris, N. and Barford, R., eds.). p. 268–284. American Chemical Society. Washington D. C.
- Lanier, T. C. 2000. Surimi gelation chemistry. *In* *Surimi and Surimi Seafood*. (Park, J. W., ed.). p. 237–265. Marcel Dekker, Inc. New York.
- Lee, C. M. 1984. Surimi process technology. *Food Technol.* 38 (11): 69–80.
- Lee, J. J., Chen, H. C. and Jiang, S. T. 1993. Purification and characterization of proteinases identified as cathepsin L and L-like (58 kDa) from mackerel (*Scomber australasicus*). *Biosci. Biotechnol. Biochem.* 57: 1470–1476.

- Lee, N. and Park, J. W. 1998. Calcium compounds to improve gel functionality of Pacific whiting and Alaska pollack surimi. *J. Food Sci.* 63: 969–974.
- Lee, J. J., Tzeng, S. S. and Jiang, S. T. 2000a. Purification and characterization of low molecular weight kininogen from pig plasma. *J. Food Sci.* 65: 81–86.
- Lee, J. J., Tzeng, S. S., Wu, J. and Jiang, S. T. 2000b. Inhibition of thermal degradation of mackerel surimi by pig plasma protein and L-kininogen. *J. Food Sci.* 65: 1124–1129.
- Lee, Y. Z., Aishima, T., Nakei, S. and Sim, J. S. 1987. Optimization for selective fractionation of bovine blood plasma proteins using poly (ethylene glycol). *J. Agric. Food Chem.* 35: 658–962.
- Leinot, A. and Cheftel, J. C. 1990. Influence of the fishing season and of chilled or frozen storage of sardine on the solubility and gelling properties of myofibrillar proteins. *In* Chilling and Freezing of New Fish Products. Paris, France. 18–20 September 1990. The International Institute of Refrigeration.
- Lin, T. S. and Lanier, T. C. 1980. Properties of alkaline protease from the skeletal muscle of Atlantic croaker. *J. Food Biochem.* 4: 17–28.
- Liu, D., Kanoh, S. and Niwa, E. 1996. Effect of cysteine protease inhibitor on the setting of Alaska pollack surimi paste. *Nippon Suisan Gakkaishi.* 62: 275–279.
- Liu, G., and Xiong, Y. L. 1997. Gelation of chicken muscle myofibrillar proteins treated with protease inhibitors and phosphates. *J. Agric. Food Chem.* 45: 3437–3442.
- Lorand, L. and Konishi, K. 1964. Activation of the fibrin stabilizing factor of plasma by thrombin. *Arch. Biochem. Biophys.* 105: 58–67.
- Lorier, M. A., Aitken, B. L. 1991. Method for treating fish with α_2 -macroglobulin. U. S. Patent 5,013,568
- Lowry, O. H., Rosebrough, N. J., Farr, L. A. and Randall, R. J. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193: 256–275.
- Lou, X., Wang, C., Xiong, Y. L., Wang, B. and Mims, S. D. 2000. Gelation characteristics of paddlefish (*Polydon spathula*) surimi under different heating conditions. *J. Food Sci.* 65: 394–398.
- Lu, G. H. and Chen, T. C. 1999. Application of egg white and plasma powders as muscle food binding agents. *J. Food Eng.* 42: 147–151.

- McCullough, J. 1995. Tranfusion medicine. *In* Blood: Principle & Practical of Hematology. (Handin, R. I., Lux, S. E. and Stossel, T. P., eds.). p. 1950–1958. Lippincott company. Philadelphia.
- Makinodan, Y., Toyohara, H. and Ikeda, S. 1984. Comparison of muscle proteinase activity among fish species. *Comp. Biochem. Physiol.* 79B: 129–134.
- Makinodan, Y., Toyohara, H. and Niwa, E. 1985. Implication of muscle alkaline proteinase in the texture degradation of fish meat gel. *J. Food Sci.* 50: 1351–1355.
- Makinodan, Y., Yamamoto, M. and Shimidu, W. 1963. Studies on muscle of aquatic animals protease in fish muscle. *Bull. Jap. Soc. Sci. Fish.* 29: 776–780.
- Mashiko, H., Miyamoto, K. and Takahashi, H. 1998. Studies on porcine high molecular weight kininogen. An improved method for the purification of porcine high molecular weight kininogen and cleavage of the kininogen by the action of porcine plasma kallikrein. *Comp. Biochem. Physiol.* 120B: 647–656.
- Mashiko, H. and Takahashi, H. 1997. Canine plasma kininogen: evidence for the presence of two kininogens and purification of high molecular weight kininogen and characterization as multi-functional molecule. *Comp. Biochem. Physiol.* 117B: 535–543.
- McFarland, R. 1990. Arrowtooth flounder protease inhibitor injection project, final report. Alaska Fisheries Development Foundation Inc., Anchorage AK, SK Coop. Agr. 88-ABH-00007.
- Morrissey, M. T., Wu, J. W., Lin, D. and An, H. 1993. Protease inhibitor effects on torsion measurements and autolysis of Pacific whiting surimi. *J. Food Sci.* 58: 1050–1054.
- Morrissey, P. A., Mulvihill, D. M. and Oriordan, D. 1991. Functional properties of blood proteins. *In* Developments in Food Proteins. Vol II. (Vtudson, B. J. F., ed.). p. 125–166. Elsevier Applied Science. London.
- Martone, C. B., Busconi, L., Folco, E. J., Trucco, R. E. and Sanchez, J. J. 1986. A simplified myosin from marine fish species. *J. Food Sci.* 51: 1554–1555.
- Masaki, T., Shimomukai, M., Miyauchi, Y., One, S., Tuchiya, T., Mastuda, T., Akazawa, H. and Soejima, M. 1993. Isolation and characterization of the protease responsible for jellification of Pacific hake muscle. *Nippon Suisan Gakkaishi.* 59: 683–690.

- Meyer, E. W. and Williams, L. D. 1976. Soy protein concentrates and isolate. *In* World Soybean Research. (Danville, I. L., ed.). p. 904. Interstate Printers and Publishers. New York.
- Montero, P. and Gomez-Guillen, C. 1996. Thermal aggregation of sardine muscle proteins during processing. *J. Agric. Food Chem.* 44: 3625-3630.
- Montero, P., Hurtado, J. L. and Perez-Mateos, M. 2000. Microstructural behavior and gelling characteristics of myosystem protein gels interacting with hydrocolloids. *Food Hydrocolloid.* 14: 455-461.
- Moure, F., Rendueles, M. and Diaz, M. 2003. Coupling process for plasma protein fractionation using ethanol precipitation and ion exchange chromatography. *Meat Sci.* 64: 391-398.
- Nakayasu, T. and Nagasawa, S. 1979. Studies on human kininogens I. Isolation, characterization, and cleavage by plasma kallikrein of high molecular weight (HMW)-kininogen. *J. Biochem.* 85: 249-258.
- Niwa, E. 1985. Functional aspect of surimi. *In* Proceedings of the International Symposium on Engineered Seafood Including Surimi. (Martin, R. E. and Collete, R. L., eds.). p. 141-147. National Fisheries Institute. Washington D.C.
- Niwa, E. 1992. Measurement of surimi composition and functional properties. *In* Surimi Technology. (Lanier, T. C. and Lee, C. M., eds.). p. 389-427. Marcel Dekker, Inc. New York.
- Niwa, E., Wang, T. T., Kanoh, S. and Nakayama, T. 1988. Contribution of gelling substance to muscular protein network structure within kamaboko. *Nippon Suisan Gakkaishi.* 94: 989-992.
- Ng, C. S. 1987. Measurement of free and expressible drips. *In* Laboratory Manual on Analytical Methods and Procedure for Fish and Fish Products. (Hasegawa, H. ed.) A4.1-A4.2. Southeast Asian Fisheries Development Center. Singapore.
- Nowsad, A. A. K. M., Kanoh, S. and Niwa, E. 1994. Setting of transglutaminase free actomyosin paste prepared from Alaska pollack surimi. *Fish. Sci.* 60, 295-297.
- Ockerman, H. W. and Hansen C. L. 2000. Animal by-product processing & utilization. Technomic publishing Co. Inc. Lancaster.
- Ogawa, M., Kanamaru, J., Miyashita, H, Tamiya, T. and Tsuchiya, T. 1995. Alpha-helical structure of fish actomyosin: changes during setting. *J. Food Sci.* 60: 297-299.

- Otlewski, J., Krowarsch, D. and Apostoluk, W. 1999. Protein inhibitors of serine proteinases. *Acta. Biochem. Pol.* 46: 531–565.
- Papa, I., Alvarez, C., Verrez-Bagnis, V., Fleurence, J. and Benyamin, Y. 1996. Post mortem release of fish white muscle α -actinin as a marker of disorganization. *J. Sci. Food Agric.* 72: 63–70.
- Pares, D. and Ledward, D. A. 2001. Emulsifying and gelling properties of porcine blood plasma as influenced by high-pressure processing. *Food Chem.* 74: 139–145.
- Pares, D., Sagure, E., Saurina, J., Sunol, J. J. and Carretero, C. 1998. Functional properties of heat induced gels from liquid and spray dried porcine blood plasma as influenced by pH. *J. Food Sci.* 63: 958–961.
- Park, J. W. 1994. Functional protein additives in surimi gels. *J. Food Sci.* 59: 525–527.
- Park, J. W. 2000. Ingredient technology and formulation development. *In* Surimi and Surimi Seafood. (Park, J. W., ed.). p. 343–392. Marcel Dekker. New York.
- Peters, T. 1975. Serum albumin. *In* The Plasma Protein. Vol. 2. (Putnam, F. W., ed.). p. 133–172. Academic Press. New York.
- Pietrasik, Z. and Li-Chan, E. C. Y. 2002. Binding and textural properties of beef gels as affected by protein, κ -carrageenan and microbial transglutaminase addition. *Food Res. Int.* 35: 91–98.
- Pomeranz Y. 1991. Functional properties of food components. Academic Press, Inc. New York:
- Porter, R. W., Koury, B. and Kudo, G. 1993. Inhibition of protease activity in muscle extracts and surimi from Pacific whiting (*Merluccius productus*), and arrowtooth flounder (*Atheresthes stomias*). *Marine Fish. Rev.* 55: 10–15.
- Putnam, F. W. 1975. Alpha, beta, gamma, omega—the structure of the plasma proteins. *In* The Plasma Proteins. Vol I. (Putnam, F. W., ed.). p. 58–125. Academic Press. New York.
- Ramos-Martinez, E., Morales-Gonzalez, O. G., Ramirez, J. A., Garcia-Carreno, F. L. and Montejano-Gaitan, J. G. 1999. Determination of the modori phenomenon and its origin in surimi of 5 fish species from the Gulf of Mexico. *Int. J. Food Sci. Technol.* 5: 397–406.

- Rawdkuen, S., Benjakul, S., Visessanguan, W. and Lanier, T. C. 2004a. Chicken plasma protein affects surimi from bigeye snapper (*Priacanthus tayenus*). *Food Hydrocolloid*. 18: 259–270.
- Rawdkuen, S., Benjakul, S., Visessanguan, W. and Lanier, T. C. 2004b. Chicken plasma protein: proteinase inhibitory activity and its effect on surimi gel properties. *Food Res. Int.* 37: 156–165.
- Rawdkuen, S., Benjakul, S., Visessanguan, W. and Lanier, T. C. 2005a. Combination effects of chicken plasma protein and setting phenomenon on gel properties and cross-linking of bigeye snapper muscle proteins. *Lebensm.-Wiss. u. Technol.* 38: 353–362.
- Rawdkuen, S., Benjakul, S., Visessanguan, W. and Lanier, T. C. 2005b. Partial purification and characterization of cysteine proteinase inhibitor from chicken plasma. *Comp. Biochem. Physiol.* submitted.
- Rawdkuen, S., Benjakul, S., Visessanguan, W. and Lanier, T. C. 2005c. Cysteine proteinase inhibitor from chicken plasma: fractionation, characterization and its autolysis inhibition of myofibrillar proteins. *Food Chem.* submitted.
- Rawdkuen, S., Benjakul, S., Visessanguan, W. and Lanier, T. C. 2005d. Fractionation and characterization of cysteine proteinase inhibitor from chicken plasma. *J. Food Biochem.* 29: 486–503.
- Record, M. T. J., Zhang, W. and Anderson, C. F. 1998. Analysis of effects of salts and uncharged solutes on protein and nucleic acid equilibria and processes: a practical guide to recognizing and interpreting polyelectric effects, Hofmeister effects and osmotic effects of salts. *Adv. Protein Chem.* 51: 281–353.
- Reppond, K. D. and Babbitt, J. K. 1993. Protease inhibitors affect physical properties of arrowtooth flounder and walleye pollock surimi. *J. Food Sci.* 58: 96–98.
- Richard, K., Owusu, A. and Kritika, M. 1999. The heat resistance and conformational plasticity of Kunitz soybean trypsin inhibitor. *J. Food Biochem.* 23: 209–224.
- Ristol, D. P., Rabaneda, G. F. and Lopez, H. M. T. 2002. Process for the production of virus-inactivated human gammaglobulin G. U. S. Patent 6,875,848.
- Ritonja, A., Krizaj, I., Mesko, P., Kopitar, M., Lucovnik, P., Strukelj, B., Pungercar, J., Buttler, D. J., Barrett, A. J. and Turk, V. 1990. The amino acid sequence of a novel inhibitor of cathepsin D from potato. *FEBS Lett.* 267: 13–15.

- Robinson, H. W. and Hodgen, C. G. 1940. The biuret reaction in the determination of serum protein. I. A study of the condition necessary for the production of the stable color which bears a quantitative relationship to the protein concentration. *J. Biol. Chem.* 135: 707–725.
- Roussel, H. and Cheftel, J. C. 1990. Mechanism of gelation of sardine protein: Influence of thermal processing and various additives on the texture and protein solubility of kamaboko gel. *Int. J. Food Sci. Technol.* 25: 260–280.
- Saeki, H., Iseya, Z., Sugiura, S. and Seki, N. 1995. Gel forming characteristics of frozen surimi from chum salmon in the presence of protease inhibitors. *J. Food Sci.* 60: 917–921, 928.
- Saito, A. and Kitano, Y. 2000. Comparative studies on the cysteine proteinase inhibitory capacity of mammalian blood. *Tohoku J. Exp. Med.* 190: 83–92.
- Sakvesen, G., Nagase, H. 1989. Inhibition of proteolytic enzymes. *In Proteolytic Enzymes a Practical Approach.* (Beynon, R. and Bond, J., eds.). p. 83–104. IRL Press. Oxford.
- Salvesen, G., Parkes, C., Abrahamson, M., Grubb, A. and Barrett, A. J. 1986. Human low- M_r kininogen contains three copies of a cystatin sequence that are divergent in structure and in inhibitory activity for cysteine proteinases. *Biochem. J.* 234: 429–434.
- Sano, T., Noguchi, S. F., Tsuchiya, T. and Matsumoto, J. J. 1988. Dynamic viscoelastic behavior of natural actomyosin and myosin during thermal gelation. *J. Food Sci.* 53: 924–928.
- Sareevoravitikul, R., Simpson, B. K. and Ramaswamy, H. 1996. Effect of crude α_2 macroglobulin on properties of bluefish (*Pomatomus saltatrix*) gels prepared by high hydrostatic pressure and heat treatment. *J. Food Biochem.* 20: 49–63.
- Seki, N., Uno, H., Lee, N. H., Kimura, I., Toyoda, K., Fujita, T. and Arai, K. 1990. Transglutaminase activity in Alaska pollack muscle and surimi and its reaction with myosin B. *Nippon Suisan Gakkaishi.* 56: 125–132.
- Seymour, T. A., Morrissey, M. T., Peters, M. Y. and An, H. 1994. Purification and characterization of Pacific whiting protease. *J. Agric. Food Chem.* 42: 2421–2427.
- Seymour, T. A., Peter, M. Y., Morrissey, M. T. and An, H. 1997. Surimi gel enhancement by bovine plasma proteins. *J. Agric. Food Chem.* 45: 2919–2923.

- Sharma, V. K. and Kalonia, D. S. 2004. Polyethylene glycol-induced precipitation of interferon alpha-2a followed by vacuum drying: development of a novel process for obtaining a dry, stable powder. *Am. Assoc. Pharm. Sci.* 6: 1-14.
- Shimizu, Y., Toyohara, H. and Lanier, T. C. 1992. Surimi production from fatty and dark fleshed fish species. *In Surimi Technology*. (Lanier, T. C. and Lee, C. M., eds.) p. 181-207. Marcel Dekker. New York.
- Sikorski, Z. E. and Kohakowski, E. 2000. Endogenous enzyme activity and seafood quality: influence of chilling, freezing, and other environmental factors. *In Seafood Enzymes*. (Haard, N.F. and Simpson, B. K., eds.) p. 451-487. Marcel Dekker, Inc. New York.
- Silva, J. G., Morais, H. A., Oliveira, A. L. and Silvestre, M. P. C. 2002. Addition effects of bovine blood globin and sodium caseinate on the characteristics of raw and cooked ham plate. *Meat Sci.* 63: 177-184.
- Steel, R. G. D. and Torrie, J. H. 1980. *Principle and Procedure of Statistics*. 2nd ed. MacGraw-Hill. New York.
- Stoknes, I. and Rustad, T. 1995. Proteolytic activity in muscle from Atlantic salmon (*Salmo salar*). *J. Food Sci.* 60: 711-714.
- Stoknes, I., Rustad, R., Mohr, V. 1993. Comparative studies of the proteolytic activity of tissue extracts from cod (*Gadus morhua*) and herring (*Clupea harengus*). *Comp. Biochem. Physiol.* 106B: 613-619.
- Stratil, A., Cizova-Schroffelova, D., Tojny, J. and Hradecky, J. 1990. Polymorphism of pig serum α -protease inhibitors-3 (PI3) and assignment of the locus to the Pi1, Po1A, Po1B, Pi2, Igh linkage group. *Animal Genet.* 21: 267-276.
- Su, H., Lin, T. S and Lanier, T. C. 1981. Investigation into potential sources of heat-stable alkaline protease in mechanically separated Atlantic croaker (*Micropogon undulatus*). *J. Food Sci.* 46: 1654-1656, 1664.
- Sugo, T., Kato, H., Iwanaga, S. and Fujii, S. 1979. Occurrence of Leu-Lys-bradykinin and histidine-rich peptide in high-molecular weight kininogen isolated from horse plasma. *Biochim. Biophys. Acta.* 579: 474-478.
- Suwansakornkul, P., Itoh, Y., Hara, S. and Obatake, A. 1993. Identification of proteolytic activities of gel-degradation factors in three lizardfish species. *Nippon Suisan Gakkaishi.* 59: 1039-1045.

- Tabilo-Munizaga, G. and Barbosa-Canovas, G. 2005. Rheology for the food industry. *J. Food Eng.* 67: 147-156.
- Tahtinen, V., Weber, E., Gunther, D., Ylonen, A., Kalkkinen, N., Olsen, R., Jarvinen, M., Soderstrom, K. O., Rinne, A., Bjorklund, H. and Bogwald, J. 2002. Immunolocalization of cysteine proteinases (cathepsins) and cysteine proteinase inhibitors (salarin and salmon kininogen) in Atlantic salmon, *Salmo salar*. *Cell Tissue Res.* 310: 213-222.
- Takahashi, N., Takahashi, Y. and Putnam, F. W. 1986. Primary structure of blood coagulation factor XIIIa (fibrinolygase, transglutaminase) from human placenta, *In Proceedings of the National Academy of Sciences of the United State of American.* 85: 5829-5833.
- Thomas, D., Richards, A. and Kay, J. 1989. Inhibition of aspartic proteinases by alpha-2-macroglobulin. *J. Biochem.* 259: 905-907.
- Timashaff, S. N. and Arakawa, T. 1997. *Protein Structure: A Practical Approach.* IRL Press. Oxford.
- Torres, M. R., Marin, F. R., Ramos, A. J. and Soriano, E. 2002. Study of operation conditions in concentration of chicken blood plasma proteins by ultrafiltration. *J. Food Eng.* 54: 215-219.
- Toyohara, H., Kinoshita, M. and Shimizu, Y. 1990a. Proteolytic degradation of threadfin bream meat gel. *J. Food Sci.* 55: 259-260.
- Toyohara, H., Sakata, T., Yamashita, K., Kinoshita, M. and Shimizu, Y. 1990b. Degradation of oval-file fish meat gel caused by myofibrillar proteinase. *J. Food Sci.* 55: 364-368.
- Toyohara, H. and Simizu, Y. 1988. Relation of the rigor mortis of fish body and the texture of the muscle. *Nippon Suisan Gakkaishi.* 54: 1795-1798.
- Traore, F. T. and Meunier, J. C. 1992. Cross-linking activity of placental FXIIIa on whey proteins and casein. *J. Agric. Food Chem.* 40: 399-402.
- Travis, J. and Salvensen, G. S. 1983. Human plasma proteins inhibitors. *Ann. Rev. Biochem.* 52: 655-709.
- Tseng, T. F., Liu, D. C. and Chen, M. T. 2000. Evaluation of transglutaminase on the quality of low salt chicken meat balls. *Meat Sci.* 55, 427-431.

- Tsukamasa, Y., Miyake, Y., Ando, M. and Makinodan, Y. 2002. Total activity of transglutaminase at various temperatures in several fish meats. *Fish. Sci.* 68: 929-933.
- Tsukamasa, Y. and Simizu, Y. 1989. The gel forming properties of the dorsal muscle from *Clupeiformes* and *Salmonoidei*. *Bull. Jap. Soc. Sci. Fish.* 55: 529-534.
- Turk, V. and Bode, W. 1991. The cystatins: protein inhibitors of cysteine proteinases. *FEBS Lett.* 285: 213-219.
- Turk, B., Turk, V. and Salvesen, G. S. 2002. Regulating cysteine protease activity: essential role of protease inhibitors as guardians and regulators. *Curr. Pharm. Des.* 8: 1623-1637.
- Turpeinen, U., Syvanen, A. C. and Iwanaga, S. 1981. HMW and LMW kininogens. *Methods Enzymol.* 80: 172-198.
- Tybor, P. T., Dill, C. W. and Landmann, W. A. 1975. Functional properties of proteins isolated from bovine by a continuous pilot process. *J. Food Sci.* 40: 155-159.
- Visessanguan, W. and An, H. 2000. Effects of proteolysis and mechanism of gel weakening in heat-induced gelation of fish myosin. *J. Agric. Food Chem.* 48: 1024-1032.
- Visessanguan, W., Benjakul, S. and An, H. 2000. Porcine plasma proteins as a surimi protease inhibitor: effects on actomyosin gelation. *J. Food Sci.* 65: 607-611.
- Visessanguan, W., Benjakul, S. and An, H. 2003. Purification and characterization of cathepsin L in arrowtooth flounder (*Atheresthes stomias*) muscle. *Comp. Biochem. Physiol.* 134B: 477-487.
- Visessanguan, W., Menino, A. R., Kim, S. M. and An, H. 2001. Cathepsin L: a predominant heat activated proteinase in arrowtooth flounder muscle. *J. Agric. Food Chem.* 49: 2633-2640.
- Wan, J., Kimura, I., Satake, M. and Seki, N. 1994. Effect of calcium ion concentration of the gelling properties and transglutaminase activity of walleye pollack surimi paste. *Fish. Sci.* 60: 107-113.
- Wasson, D. H., Babbitt, J. K. and French, J. S. 1992a. Characterization of a heat stable protease from arrowtooth flounder: *Atheresthes stomias*. *J. Aquat. Food Prod. Technol.* 1 (3/4): 167-182.

- Wasson, D. H., Reppond, K. D., Babbitt, J. K. and French, J. S. 1992b. Effects of additives on proteolytic and functional properties of arrowtooth flounder surimi. *J. Aquat. Food Prod. Technol.* 1(3/4): 147-165.
- Weerasinghe, V. C., Morrissey, M. T., and An, H. 1996a. Characterization of active components in food-grade protease inhibitors for surimi manufacture. *J. Agric. Food Chem.* 44: 2584-2590.
- Weerasinghe, V. C., Morrissey, M. T., Chung, Y. and An, H. 1996b. Whey protein concentrates as a proteinase inhibitor in Pacific whiting surimi. *J. Food Sci.* 61: 367-371.
- White, A., Handler, P. and Smith, E. 1973. *Principles of Biochemistry*. 5th Ed. McGraw Hill Book Company. New York.
- Whitaker, J. R. 1994. *Principles of Enzymology for the Food Sciences*. 2nd Ed. Marcel Dekker, Inc. New York.
- Wu, S. M. and Pizzo, S. V. 1999. Mechanism of hypochlorite-mediated inactivation of proteinase inhibition by α_2 -macroglobulin. *Biochemistry*. 38: 13983-13990.
- Xiong, Y. L. 1993. A comparison of the rheological characteristics of different fractions of chicken myofibrillar proteins. *J. Food Biochem.* 16: 217-227.
- Xiong, Y. L. 1997. Structure-function relationships of muscle proteins. *In Food Proteins and Their Applications*. (Damodaran, S. and Paref, A., eds.). p. 341-392. Dekker. New York.
- Xiong, Y. L. 1997. Structure-function relationships of muscle proteins: viscoelastic changes related to heating procedures. *J. Food Sci.* 59: 734-738.
- Yamamoto, T. 1987. Characterization of guinea pig high-molecular weight kininogen as multi-functional molecule. *Biochim. Biophys. Acta.* 914: 259-274.
- Yamashita, M. and Konagaya, S. 1990. Participation of cathepsin L into extensive softening of the muscle of chum salmon caught during spawning migration. *Nippon Suisan Gakkaishi.* 56: 1271-1277.
- Ylonen, A., Helin, J., Bogwald, J., Jaakola, A., Rinne, A. and Kalkkinen, N. 2002. Purification and characterization of novel kininogens from spotted wolffish and Atlantic cod. *Eur. J. Biochem.* 269: 2639-2646.
- Ylonen, A., Rinne, A., Herttuainen, J., Bogwald, J., Jarvinen, M. and Kalkkinen, N. 1999. Atlantic salmon (*Salmo salar* L.) skin contains a novel kininogen and another cysteine proteinase inhibitor. *Eur. J. Biochem.* 266: 1066-1072.

- Yoon, W. B., Kim, B. Y. and Park, J. W. 1999. Rheological characteristics of fibrinogen-thrombin solution and its effects on surimi gels. *J. Food Sci.* 64: 291-294.
- Yoshida, K., Saito, A. and Sinohara, H. 1989. Isolation and partial characterization of a novel form of low molecular weight kininogen from guinea pig plasma. *Biochem. Int.* 19: 1339-1344.
- Zeece, G. M., Woods, T. L., Keen, M. A. and Reville, W. J. 1992. Role of endogenous and inhibitors in postmortem muscle protein degradation. *Proc. Recip. Meat Conf.* 45: 51-61.
- Ziegler, G. R. and Acton, J. C. 1984. Mechanisms of gel formation by proteins of muscle tissues. *Food Technol.* 38 (5): 77-80, 82.
- Ziegler, G. R. and Foegeding, E. A. 1991. The gelation of proteins. *In Advances in Food and Nutrition Research.* (Kinsella, J. E., ed.). p. 203-298. Academic Press. New York.