

CONTENTS

	Page
Contents	vii
List of tables	ix
List of figures	xi
Chapter 1. Introduction	1
Review of literature	4
The characteristics of the genus <i>Padina</i>	4
Diversity of <i>Padina</i> in Thailand	5
Growth and reproduction of <i>Padina</i> species	6
Recruitment	8
Environmental factors	9
Hypothesis	11
Objectives	11
Research questions	12
Chapter 2. Materials and Methods	13
Study sites	13
Growth study	15
Reproduction study	16
Recruitment study	21
Statistical analyses	23

CONTENTS (Continued)

	Page
Chapter 3. Results	24
Size structure and reproductive potential	24
The reproductive cycle and the time of production of new individuals	32
Recruitment study	40
Chapter 4. Discussion	57
Life phase of <i>P. boryana</i>	57
Population structure, reproductive potential and recruitment of <i>P. boryana</i>	59
Chapter 5. Conclusions	63
References	64
Appendix	74
Vitae	86

LIST OF TABLES

Table	Page
1. The average number of each type of sporangium arrangement.	20
2. A comparison of mean surface area of <i>P. boryana</i> blade at Sirinart National Park and Tang Khen Bay from September 2005 to August 2006	29
3. A comparison of size class distribution at Sirinart National Park and Tang Khen Bay from September 2005 to August 2006.	30
4. The average number of released spores at Sirinart National Park and Tang Khen Bay from September 2005 to August 2006.	31
5. The average percentage of reproductive stages at Sirinart National Park and Tang Khen Bay.	34
6. Summary of percentages of each life stage varied throughout the year, from September 2005 to August 2006 at Sirinart National Park and Tang Khen Bay.	35
7. The average of percentage of maturation stages at Sirinart National Park and Tang Khen Bay.	37
8. The average number of spores produced at Sirinart National Park and Tang Khen Bay from September 2005 to August 2006.	39
9. A comparison of the percentage of filamentous greens, filamentous reds and <i>P. boryana</i> at all levels in the permanent plots at Sirinart National Park from October 2005 to August 2006.	44

LIST OF TABLES (Continued)

Table	Page
10. A comparison of the percentage of filamentous greens, filamentous reds and <i>P. boryana</i> at all levels in the permanent plots at Tang Khen Bay from October 2005 to August 2006.	45
11. The average percentage of <i>P. boryana</i> recruitment at all shore levels at Sirinart National Park and Tang Khen Bay from October 2005 to August 2006.	46
12. The average length (cm.) of <i>P. boryana</i> in the permanent plots at Sirinart National Park and Tang Khen Bay from October 2005 to August 2006; D: <i>Dictyosphaeridia</i> stage found.	47
13. The correlation coefficient value of physio-chemical parameters and biological characteristics of <i>P. boryana</i> .	51

LIST OF FIGURES

Figure	Page
1. The whole thallus of <i>Padina boryana</i>	5
2. Diagram of the life cycle of <i>Padina boryana</i> (modified from Lee, 1999).	7
3. The longitudinal cross-section of <i>Padina boryana</i> (X400) showing the oogonia packed on the blade.	8
4. The longitudinal cross-section of <i>Padina boryana</i> (X100) showing the reproductive sori of antheridia.	8
5. Map of Phuket showing the two study locations; Sirinart National Park and Tang Khen Bay.	14
6. Populations of <i>Padina boryana</i> Thivy from study sites; SNP (left) and TKB (right).	14
7. The whole thallus of <i>P. boryana</i> showing the length and diameter measurements.	16
8. Sporangium arrangement on the surface of the blade (X40); A: type 1, B: type 2, C: type 3. D: type 4, E: type 5.	19
9. A one centimeter length of <i>P. boryana</i> reproductive line was randomly selected to investigate the number of sporangia/sorus.	20
10. System of collection to assess the population distribution, reproduction and recruitment of <i>P. boryana</i> .	22

LIST OF FIGURES (Continued)

Figure	Page
11. Size class of <i>P. boryana</i> populations from September 2005 to August 2006 comparing Sirinart National Park and Tang Khen Bay.	26
12. Comparison of the average number of released spores/thalli from September 2005 to August 2006 at Sirinart National Park and Tang Khen Bay.	31
13. Comparison of percentages of life phases of <i>P. boryana</i> from September 2005 to August 2006 at Sirinart National Park and Tang Khen Bay.	34
14. Comparison of the percentages of mature (stages 3-5) thalli from September 2005 to August 2006 at Sirinart National Park and Tang Khen Bay.	36
15. Comparison of the average number of spores/thalli from September 2005 to August 2006 at Sirinart National Park and Tang Khen Bay.	38
16. Percentage cover of <i>P. boryana</i> from October 2005 to August 2006 at Sirinart National Park.	48
17. Percentage cover of <i>P. boryana</i> from October 2005 to August 2006 at Tang Khen Bay.	49
18. Comparison of seawater temperatures from September 2005 to August 2006 at Sirinart National Park and Tang Khen Bay.	50

LIST OF FIGURES (Continued)

Figure	Page
19. Comparison of air temperatures from September 2005 to August 2006 at Sirinart National Park and Tang Khen Bay.	50
20. Comparison of salinity (ppt) from September 2005 to August 2006 at Sirinart National Park and Tang Khen Bay.	51
21. The correlation of average surface areas and air temperatures from September 2005 to August 2006 at Sirinart National Park.	52
22. The correlation of average percentages of female gametophytes and air temperature from September 2005 to August 2006 at Tang Khen Bay.	52
23. The correlation of average percentages of female gametophytes and water temperature from September 2005 to August 2006 at Tang Khen Bay.	53
24. The correlation of average percentages of male gametophytes and salinity from September 2005 to August 2006 at Tang Khen Bay.	53
25. The correlation of average percentages of sporophytes and salinity from September 2005 to August 2006 at Tang Khen Bay.	54
26. The correlation of average percentages of mature sporophytes and salinity from September 2005 to August 2006 at Tang Khen Bay.	54
27. The correlation of average percentages of reproduction and salinity from September 2005 to August 2006 at Tang Khen Bay.	55

LIST OF FIGURES (Continued)

Figure		Page
28.	The correlation of average numbers of spores produced and salinity from September 2005 to August 2006 at Tang Khen Bay.	55
29.	The correlation of average numbers of spores released and salinity form September 2005 to August 2006 at Tang Khen Bay.	56
30.	The correlation of average surface areas and salinity from September 2005 to August 2006 at Tang Khen Bay.	56