

References

- Al-Saikhan, M.S., Howard, L.R. and Miller, J.R. 1995. Antioxidant activity and total phenolics in different genotypes of potato (*Solanum tuberosum*, L.). *J. Food Sci.* 60: 341-343.
- Aruoma, O.I. 1994. Deoxyribose assays for detecting hydroxyl radicals. *Methods Enzymol.* 233: 57-66.
- Aruoma, O.I., Spencer, J.P.E., Rossi, R., Aeschbach, A.K., Mahmood, N., Munoz, A., Murcia, A., Butler, J. and Halliwell, B. 1996. An evaluation of the antioxidant and antiviral action of extracts of rosemary and provençal herbs. *Food Chem. Toxicol.* 34: 449-456.
- Arts, I.C.W., van de Putte, B. and Hollman, P.C.H. 2000. Catechin contents of foods commonly consumed in the Netherlands. *J. Agric. Food Chem.* 48: 1752-1757.
- Asakawa, T., Matsushita, S. 1980. Coloring condition of thiobarbituric acid test for detection lipid hydroperoxides. *Lipids.* 15: 137-140.
- Asano, N. 2001. Polyhydroxylated alkaloids isolated from mulberry trees (*Morus alba* L.) and silkworms (*Bombyx mori* L.). *J. Agric. Food Chem.* 49: 4208-4213.
- Bandarra, N.M., Campos, R.M., Batista, I., Nunes, M.L. and Empis, J.M. 1999. Antioxidant synergy of α -tocopherol and phospholipids. *J. Am. Oil Chem. Soc.* 76: 905-913.
- Barton, G.M., Evan, R.S. and Gardner, J.A.F. 1952. Paper chromatography of phenolic substances. *Nature.* 17: 249-252.
- Baublis, A., Decker, E.A. and Clydesdale, F.M. 2000. Antioxidant effect of aqueous extracts from wheat based ready-to-eat breakfast cereals. *Food Chem.* 68: 1-6.
- Beecher, G.R. 1999. Flavonoids in Foods. *In* Antioxidant Food Supplements in Human Health. (Packer, L., Hiramatsu, M. and Yoshikawa, T., eds.). p. 269-281. Academic Press. London.

- Benavente-García, O., Castillo, J., Lorente, J., Ortuño, A. and Del-Rio, J.A. 2000. Antioxidant activity of phenolics extracted from *Olea europaea* L. leaves. *Food Chem.* 68: 457-462.
- Benzie, I.F.F. and Szeto, Y.T. 1999. Total antioxidant capacity of teas by the ferric reducing antioxidant power assay. *J. Agric. Food Chem.* 47: 633-636.
- Bhatia, I. and Ullah, M. 1968. Qualitative and quantitative study of polyphenols of different organs and some cultivated varieties of tea plant. *J. Sci. Food. Agri.* 19: 535.
- Blois, M.S. 1958. Antioxidant determination by use of a stable free radical. *Nature.* 4617: 1198-1200.
- Bors, W., Hellers, W., Michel, C. and Saran, M. 1990. Flavonoids as Antioxidants: Determination of Radical-Scavenging Efficiencies. *In Methods in Enzymology.* (Packer, L. and Glazer, A.N., eds.). p. 343-355. Academic Press. San Diego.
- Brand-Williams, W., Cuvelier, M.E. and Berset, C. 1995. Use of a free radical method to evaluate antioxidant activity. *Food Sci. Tech.* 28: 25-30.
- Brown, J.E., Khodr, H., Hider, R.C. and Rice-Evans, C.A. 1998. Structural dependence of flavonoid interactions with Cu^{2+} ions: implications for their antioxidant properties. *J. Biochem.* 330: 1173.
- Cabelli, D.E. and Bielski, B.H.J. 1983. Kinetics and mechanism for the oxidation of ascorbic acid/ascorbate by HO_2/O_2 - radicals, a pulse radiolysis and stopped-flow photolysis study. *J. Phys. Chem.* 87: 1809-1812.
- Cao, G., Sofic, E. and Prior, R.L. 1997. Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships. *Free Rad. Biol. Med.* 22: 749-760.
- Chang, S.S., Matijasevic, B.O., Hsieh, O.A.L. and Huang, C.L. 1977. Natural antioxidants from rosemary and sage. *J. Food Sci.* 42: 1102-1106.
- Cheah, P.B. and Gan, S.P. 2000. Antioxidative/antimicrobial effects of galangal and α -tocopherol in minced beef. *J. Food Protection.* 63: 404-407.
- Chen, B.H. and Tang, Y.C. 1998. Processing and stability of carotenoid power from carrot pulp waste. *J. Agric. Food Chem.* 46: 2312-2318.

- Chen, C.W. and Ho, C.T. 1995. Antioxidant properties of polyphenols extracted from green and black teas. *J. Food Lipid.* 2: 35-46.
- Chen, F. 1995. Hypoglycemic activity and mechanisms of extracts from mulberry leaves (*folium mori*) and cortex mori radicals in streptozotocin-induced diabetic mice. *Yakugaku Zasshi.* 115: 476-482.
- Chen, Z.Y. and Chan, P.T. 1996. Antioxidative activity of green tea catechins on canola oil. *Chem. Phys Lipid.* 82: 163-172.
- Chimi, H., Chillard, J., Chillard P. and Rahmani, M. 1991. Peroxyl and hydroxyl radical scavenging activity of some natural phenolic antioxidants. *J. Am. Oil. Chem. Soc.* 68: 307-312.
- Chipault, J.R., Mizuno, G.R. and Lundberg, W.O. 1995. Antioxidant properties of spices in oil-in-water emulsions. *Food Res. Intern.* 20: 443-448.
- Cho, S.Y., Miyashita, K., Miyazawa, T. Fujimoto. and Kaneda, T. 1987. Autoxidation of ethyl eicosapentaenoate and docosahexaenoate. *J. Am. Oil Chem. Soc.* 64: 876-879.
- Choi, H.S., Song, H.S., Ukeda, H. and Sawamura, M. 2000. Radical-scavenging activities of citrus essential oils and their components: detection using 1,1-diphenyl-2-picrylhydrazyl. *J. Agric. Food Chem.* 48: 4156-4161.
- Chu, D.C. 1997. Green Tea-its Cultivation, Processing of Tea Leaves for Drinking Material, and Kinds of Green Tea. *In Chemistry and Applications of Green Tea.* (Yamamoto, T., Juneja, L.R., Chu, D.C. and Kim, M., eds.). p. 1-12. Boca Raton. New York.
- Cilliers, J.J.L, and Singleton, V.L. 1990. Antioxidative phenolic ring opening under alkaline conditions as a model for natyral polyphenols in food. *J. Agric. Food Chem.* 38: 1797-1798.
- Cilliers, J.J.L, and Singleton, V.L. 1991. Characterization of the products of nonenzymic autoxidative phenolic reaction in a caffeic acid model system. *J. Agric. Food Chem.* 31: 1298-1303.

- Conde, E., Cadahia, E., Garcia-Vallejo, M. and Fernandez de Simon, B. 1998. Polyphenolic composition of *Quercus suber* cork from different Spanish provenances. *J. Agric. Food Chem.* 44: 3166-3171.
- Cort, W.M. 1982. Antioxidant properties of ascorbic acid in foods. *Adv. Chem Ser.* 200: 533-550.
- Cox, D.A. and Cohen M.L. 1996. Effects of oxidized low-density lipoprotein on vascular contraction and relaxation: clinical and pharmacological implications in atherosclerosis. *Pharmacol. Rev.* 48: 3-19.
- Cuvelier, M.E., Richard, H. and Berset, C. 1996. Antioxidative activity and phenolic composition of pilot-plant and commercial extracts of sage and rosemary. *J. Am. Oil Chem. Soc.* 73: 645-652.
- Diplock, A.T. 1992. The role of antioxidant nutrients in disease: health and nutrition. *INFORM.* 3: 1214-1217.
- Doba, T., Burton, G.W. and Ingold, K.U. 1985. Antioxidant and coantioxidant activity of vitamin C, the effect of vitamin C, ether alone or in the presence of vitamin E or a water soluble vitamin E analogue upon the peroxidation of aqueous multilamellar phospholipids liposome. *Biochem. Biophys. Acta.* 835: 293-303.
- Doi, K., Kojima, T. and Fujimoto, Y. 2000. Mulberry leaf extract inhibits the oxidative modification of rabbit and human low density lipoprotein. *Biol. Pharm. Bull.* 23: 1066-1071.
- Duh, P.D. 1998. Antioxidant activity of burdock (*Arctium lappa* Linne'): its scavenging effect on free-radical and active oxygen. *J. Am. Oil Chem. Soc.* 75: 455-461.
- Duh, P.D. and Yen, G.C. 1997. Antioxidative activity of three herbal water extracts. *Food Chem.* 60: 639-645.
- Duh, P.D., Yen, W.J., Du, P.C. and Yen, G.C. 1997. Antioxidant activity of mung bean hulls. *J. Am. Oil Chem. Soc.* 74: 1059-1063.
- Durrington, P.N. 1995. Lipoprotein and Their Metabolism. *In* Hyperlipidaemia Diagnosis and Management. p. 25-71. Butterworth-Heinemann. Oxford.

- Dziezak, J.D. 1986. Antioxidants: the ultimate answer to oxidation. *Food Technol.* 40: 94-102.
- Economou, K.D., Oreopoulou, V. and Thomopoulos, C.D. 1991. Antioxidant activity of some plant extracts of the family *Labiatae*. *J. Am. Oil Chem. Soc.* 68: 109-115.
- Eichner, K. 1981. Antioxidative effect of maillard reaction intermediates. *Prog. Food Nutr. Sci.* 5: 441-451.
- Endo, Y., Usuki, R. and Kaneda, T. 1985. Antioxidant effects on chlorophylls and pheophytin on the autoxidation of oils in the dark II. *J. Am. Oil Chem. Soc.* 62: 1387-1390.
- Esaki, H., Kawakishi, S., Morimitsu, Y. and Osawa, T. 1999. New potent antioxidative o-dihydroxyisoflavones in fermented Japanese soybean products. *Biosci. Biotechnol. Biochem.* 63: 1637-1639.
- Esterbauer, H. and K.H. Cheeseman, 1990. Determination of aldehyde lipid peroxidation products: malonaldehyde and 4-hydroxynonenal. *Methods. Enzymol.* 186: 407-420.
- Fejes, S., Blazovics, A., Lugasi, A., Lemberkovics, E., Petri, G. and Kery, A. 2000. In vitro antioxidant activity of *Anthriscus cerefolium* L. (Hoffm.) extracts. *J. Ethnopharmacol.* 69: 259-265.
- Fernandez de Simon, B., Cadahia, E., Conde, E. and Garcia-Vallejo, M.C. 1996. Low molecular weight phenolic compounds in Spanish oakwoods. *J. Agric. Food Chem.* 44: 1507-1511.
- Fogliano, V., Verde, V., Randazzo, G. and Ritieni, A. 1999. Method for measuring antioxidant activity and its application to monitoring the antioxidant capacity of wines. *J. Agric. Food Chem.* 47: 1035-1040.
- Frankel, E.N. 1980. Lipid oxidation. *Progress lipid res.* 19: 1-22.
- Frankel, E.N. 1996. Antioxidants in lipid foods and their impact on food quality. *Food Chem.* 57: 51-55.

- Frankel, E.N., Kanner, J., German, J.B., Parks, E. and Kinsella, J.E. 1993. Inhibition of oxidation of human low-density lipoprotein by phenolic substances in red wine. *Lancet*. 341: 454.
- Fuhrman, B., Lavy, A. and Aviram, M. 1995. Consumption of red wine with meals reduces the susceptibility of human plasma and low-density lipoprotein to lipid peroxidation. *Am. J. Clin. Nutr.* 61: 549-554.
- Fukatsu, S. 1992. Tea manufacturing process. *Farming Japan*. 26: 14-19.
- Gabrielska, J., Oszmianski, J. and Lamer-Zarawska, E. 1997. Protective effect of plant flavonoids on the oxidation of lecithin liposomes. *Pharmazie*. 52: 2-3.
- Gamez, E.J.C., Luyengi, L., Lee, S.K., Zhu, L.F. and Zhou, B.N. 1998. Antioxidant flavonoid glycosides from *Daphniphyllum calycinum*. *J. Nat. Prod.* 61: 706-708.
- Ghiselli, Andrea., Nardini, M., Baldi, A. and Scaccini, C. 1998. Antioxidant activity of different phenolic fractions separated from an Italian red wine. *Am. Oil Chem. Soc.* 46: 361-367.
- Giese, J. 1996. Antioxidants: tool for preventing lipid oxidation. *Food Technol.* 50: 73-80.
- Gordon, M.H. 1990. The Mechanism of Antioxidant Action *in vitro*. In *Food Antioxidants*. (Hudson, B.J.F., ed.). p. 1-18. Elsevier. London/Newyork.
- Graham, G. 1992. Green tea composition, consumption and polyphenol chemistry. *Prevent. Med.* 21: 334-350.
- Guo, Q., Zha, B., Li, M., Shen, S. and Xin, W. 1996. Studies on protective mechanisms of four components of green tea polyphenols against lipid peroxidation in synaptosomes. *Biochim. Bio-phys. Acta.* 1304: 210-222.
- Gutfinger, T. 1981. Polyphenols in olive oils. *J. Am. Oil Chem. Soc.* 58: 966-968.
- Hagerman, A.E., Riedl, K.M., Jones, G.A., Sovik, K.N., Ritchard, N.T., Hartzfeld, P.W. and Riechel, T.L. 1998. High molecular weight plant polyphenolics (tannins) as biological antioxidants. *J. Agric. Food Chem.* 46: 1887-1892.
- Halliwell, B., Gutteridge, J.M. and Cross, C.E. 1992. Free radicals antioxidants and human disease; where are we now. *J. Lab. Clin. Med.* 119: 598-620.

- Hamama, A.A. and Nawar, W.W. 1991. Thermal decomposition of some phenolic antioxidants. *J. Agric. Food Chem.* 39: 1063-1069.
- Han, D., Yi, O.S. and Shin, H.K. 1990. Antioxidative effect of ascorbic acid solubilized in oils via reversed micelles. *J. Food Sci.* 55: 247-249.
- Hara, Y. 1994. Prophylactic Functions of Tea Polyphenols. *In Food Phytochemicals for Cancer Prevention II-Tea, Spices and Herbs.* (Ho, C.T., Osawa, T., Huang, M.T. and Rosen, R.T., eds.). p. 34-50. American Chemical Society. Washington.
- He, Y. and Shahidi, F. 1997. Antioxidant activity of green tea and its catechins in a fish meat model system. *J. Agric. Food Sci.* 45: 39-45.
- Heinonen, I.M., Meyer, A.S. and Frankel, E.N. 1998. Antioxidant activity of berry phenolics on human low-density lipoprotein and liposome oxidation. *J. Agric. Food Chem.* 46: 4107-4112.
- Huang, H.C., Lee, C.R., Lee, C.P.D., Chen, C.C. and Chu, S.H. 1991. Vasorelaxant effect of emodin, an anthraquinone from Chinese herb. *J. Pharm.* 205: 289-294.
- Huang, S.W., Frankel, E.N. and German, J.B. 1994. Antioxidant activity of α - and γ -tocopherols in bulk oils and in oil-in-water emulsions. *J. Agric. Food Chem.* 42: 2108-2114.
- Hudson, B.J.F. and Mahgoub, S.E.O. 1981. Synergism between phospholipids and naturally occurring antioxidants in leaf lipids. *J. Sci. Food Agri.* 32: 208-210.
- Imida, K., Fukushima, S., Shirai, T., Ohtani, M., Nakanishi, K. and Ito, N. 1983. Promoting activities of butylated hydroxyanisole and butylated hydroxytoluene on 2-stage urinary bladder carcinogenesis and inhibition of γ -glutamyl transpeptidase positive loci development in the liver of rats. *Carcinogenesis.* 4: 895-899.
- Ibáñez, E., Oca, A., de Murga, G., López-Sebastian, S., Tabera, J. and Reglero, G. 1999. Supercritical fluid extraction and fractionation of different preprocessed rosemary plants. *J. Agric. Food Chem.* 47: 1400-1404.
- IUPAC. 1979. *Standard Methods for the Analysis of Oils, Fat and Derivatives.* 6th ed. Part I. Pergamon Press. Paris.

- Jadhav, S.J., Nimbalkar, S.S., Kulkarni, A.D. and Madhavi, D.L. 1996. Lipid Oxidation in Biological and Food Systems. *In Food Antioxidants*. (Madhavi, D.L., Deshpande, S.S. and Sulunkhe, D.K., eds.). p. 5-64. Marcel Dekker. New York.
- Julkunen-Tiito, R. 1985. Phenolic constituents in the leaves of northern willows, methods for the analysis of certain phenolics. *J. Agric. Food Chem.* 33: 213-217.
- Kahkonen, M.P., Hopia, A.I., Vuorela, H.J., Rauha, J.P., Pihlaja, K., Kujala, T.S. and Heininen, M. 1999. Antioxidant activity of plant extracts containing phenolic compounds. *J. Agric. Food Chem.* 47: 3954-3962.
- Kallithraka, S.C., Bridle, G.V.P. and Bakker, J. 1995. Survey of solvents for the extraction of grape seed phenolics. *Phytochem. Anal.* 6: 26-267.
- Kalyanaraman, B. 1995. Free Radical Mechanisms of Oxidative Modification of Low Density Lipoprotein (or the Rancidity of Body Fat). *In Oxidative Stress and Antioxidant Defenses in Biology*. (Ahmad, S., ed.). p. 96-114. Chapman & Hall. New York.
- Kashima, M., Cha, G.S., Isoda, Y., Hirano, J. and Miyazawa, T. 1991. The antioxidant effects of phospholipids on perilla oil. *J. Am. Oil Chem. Soc.* 68: 119-122.
- Kaur, H. and Perkins, J. 1991. The Free Radical-Chemistry of Food Additives. *In Free Radical and Food Additives*. (Aruoma, O.I. and Halliwell, B., ed.). p. 17-35. London
- Kawakami, S., Mizuni, M. and Tsuchida, H. 2000. Comparison of antioxidant enzyme activities between *Solanum tuberosum* L. cultivars danshaku and kitaakari during low-temperature storage. *J. Agric. Food Chem.* 48: 2117-2121.
- Khanna, S.K., Viswanatham, P.N., Krishnam, D.S. and Sawai, G.G. 1968. Extraction of total phenolics in the presence of reducing agents. *Phytochemistry*. 7: 1513-1518.
- Kim, S.Y., Gao, J.J. and Kang, H.K. 2000. Two flavonoids from the leaves of *Morus alba* induced differentiation of the human promyelocytic leukemia (HL-60) cell line. *Biol. Pharm. Bull.* 23: 451-455.

- Kim, S.Y., Kim, J.H., Kim, S.K., Oh, M.J. and Jung, M.Y. 1994. Antioxidant activities of selected oriental herb extracts. *J. Am. Oil Chem. Soc.* 71: 633-640.
- King, M.F., Boyd, L.C. and Sheldon, B.W. 1992. Antioxidant properties on Individual phospholipids in a salmon oil model system. *J. Am. Oil Chem. Soc.* 69: 545-551.
- Kleinveld, H.A., Hak-Lemmers, H.L.M., Stalenhoef, A.F.H. and Demacker, P.N.M. 1992. Improved measurement of Low-Density-Lipoprotein susceptibility to copper-induced oxidation: application of short procedure for isolating Low-Density Lipoprotein. *Clin. Chem.* 38: (10): 2066-2072.
- Koga, T., Moto, K., Nakamori, K., Yamakoshi, J., Hosoyama, H., Kataoka, S. and Ariga, T. 1999. Increase of antioxidative potential of rat plasma by oral administration of proanthocyanidin-rich extract from grape seeds. *J. Agric. Food Chem.* 47: 1892-1897.
- Koidzumi, G. 1917. Taxonomical discussion on *Morus* plants. *Bull. Imp. Sericult. Exp. Stat.* 3: 1-12.
- Kojima, Y. 2001. Retardation of the development of diabetes mellitus in murine model type 1 diabetes (NOD mouse) fed *Morus alba* leaves. *J. Jpn. Soc. Nutri. Food Sci.* 54: 361-364.
- Koketsu, M. 1997. Antioxidative Activity of Tea Polyphenols. *In Chemistry and Applications of Green Tea.* (Yamamoto, T., Juneja, L.R., Chu, D.C. and Kim, M., eds.). p. 37-50. Boca Raton. New York.
- Kono, S., Shinchi, K., Ikada, N., Yanai, F. and Imanishi, K. 1992. Green tea consumption and serum lipid profiles: a cross-sectional study in northern Kyushu, Japan. *Prev. Med.* 21: 526-531.
- Kurilich, A.C. and Juvik, J.A. 1999. Quantification of carotenoid and tocopherol antioxidants in *Zea mays*. *J. Agric. Food Chem.* 47: 1948-1955.
- Labuza, T.D. 1971. Kinetics of lipid oxidation in Foods. *Crit. Rev. Food Technol.* 7: 355-395.
- Lagouri, V. and Boskou, D. 1996. Nutrient antioxidants in oregano. *Intern. J. Food Sci. Nutr.* 47: 439-497.

- Lai, L.S., Chou, S.T. and Chao, W.W. 2001. Studies on the antioxidative activities of hsian-tsaio (*Mesona procumbens* Hemsl) leaf gum. *J. Agric. Food Chem.* 49: 963-968.
- Larrauri, J.A., Ruperez, P. and Saura-Calixto, F. 1997. Effect of drying temperature on the stability of polyphenols and antioxidant activity of red grape pomace peels. *J. Agric. Food Chem.* 44: 1390-1393.
- Lee, Y.B., Kim, Y.S. and Ashmore, C.R. 1986. Antioxidant property in ginger rhizome and its application to meat products. *J. Food Sci.* 51: 20-23.
- Leenen, R., Roodenburg, A.J.C., Tijburg, L.B.M. and Wiseman, S.A. 2000. A single dose of tea with or without milk increases plasma antioxidant activity in humans. *Eur. J. Clin. Nutr.* 54: 87.
- Lehtinen, P. and Laakso, S. 1998. Effect of extraction conditions on the recovery and potency of antioxidants in oat fiber. *J. Agric. Food Chem.* 46: 4842-4845.
- Liao, K.L. and Yin, M.C. 2000. Individual and combined antioxidant effects of seven phenolic agents in human erythrocyte membrane ghosts and phosphatidylcholine liposome systems: importance of the partition coefficient. *J. Agric. Food Chem.* 48: 2266-2270.
- Lin, Y.L., Juan, I.M., Chen, Y.L., Liang, Y.C. and Lin, J.K. 1996. Composition of polyphenols in fresh tea leaves and associations of their oxygen-radical-absorbing capacity with antiproliferative actions in fibroblast cells. *J. Agric. Food Chem.* 44: 1387-1394.
- Lissi, E.A., Modak, B., Torres, R. and Urzua, A. 1999. Total antioxidant potential of resinous exudates from *Heliotropium* species and a comparison of the ABTS and DPPH methods. *Free Rad. Res.* 30: 471-477.
- Liu, Z.Q., Ma, L.P., Zhou, B., Yang, L. and Liu, Z.L. 2000. Antioxidative effects of green tea polyphenols on free radical initiated and photosensitized peroxidation of human low density lipoprotein. *Chemistry and Physics of Lipids.* 106: 53-63.
- Lotito, S.B. and Fraga, C.G. 1998. (+)-Catechin prevents human plasma oxidation. *Free Radic. Biol. Med.* 24: 435.

- Maillard, M.N. and Berset, C. 1995. Evolution of antioxidant activity during kilning, role of insoluble bound phenolic acids of barley and malt. *J. Agric. Food Chem.* 43: 1789-1793.
- Maillard, M.N., Soum, M.H., Boivin, P. and Berset, C. 1996. Antioxidative activity of barley and malt: relationship with phenolic content. *Lebensm. Wiss.-Technol.* 29: 238-244.
- Marco, G.T. 1968. A rapid method for evaluation of antioxidants. *J. Am. Oil Chem. Soc.* 75: 1453-1455.
- Marinova, E.M. and Yanishlieva, N.V.I. 1997. Antioxidative activity of extracts from selected species of the family *Lamiaceae* in sun flower oil. *Food Chem.* 58: 245-248.
- Markwell, M.A.K., Hass, S.M., Bieber, L.L. and Tolbert, N.E. 1978. A modification of the lowry procedure to simplify protein determination in membrane and lipoprotein samples. *Anal. Bioc.* 87: 206-210.
- Masaki, H., Sakaki, S., Atsumi, T. and Sukarai, H. 1995. Active-oxygen scavenging activity of plant extracts. *Biol. Pharm. Bull.* 18: 162-166.
- Mensink, R.P., Temme, E.H.M. and Plat, J. 1998. Dietary Fats and Coronary Heart Disease. *In Food Lipids.* (Akok, C.C. and Min, D.B., eds). p. 507-523. New York.
- Meyer, A.S., Heinonen, M. and Frankel, E.N. 1998. Antioxidant interactions of catechin, cyaniding, caffeic acid, quercetin, and ellagic acid on human LDL oxidation. *Food Chem.* 61: 71-75.
- Miller, N.J., Castelluccio, C., Tijburg, L. and Rice-Evans, C. 1996. The antioxidant properties of theaflavins and their gallates esters-radical scavengers or metal chelators. *FEBS Letters.* 392: 40-44
- Milovanovic, M., Picuric, K., Vucelic, B. and Vabaski, Z. 1998. Antioxidant effects of flavonoid of *Anthriscus sylvestris* in lard. *J. Am. Oil Chem. Soc.* 73: 773-776.
- Mitsumoto, M., Faustman, C., Cassens, R.G., Arnold, R.N., Schaefer, D.M. and Scheller, K.K. 1991. Vitamin E and C improve pigment and lipid stability in ground beef. *J. Food Sci.* 56: 194-197.

- Miura, Y., Chiba, T. and Miura, S. 2000. Green tea polyphenols (flavan 3-ols) prevent oxidative modification of low density lipoproteins: an *ex vivo* study in humans. *J. Nutr. Biochem.* 11: 216.
- Miura, S., Watanabe, J., Tomita, T., Sano, M. and Tomita, I. 1995. Effect of various natural antioxidants on the Cu^{2+} - mediated oxidative modification of low density lipoprotein. *Biol. Pharmaceut. Bull.* 18: 1-4.
- Moreno, C.S., Larrauri, J.A. and Saura-Calixto, F. 1998. A procedure to measure the antiradical efficiency of polyphenols. *J. Sci. Food Agric.* 76: 270-276.
- Moure, A., Cruz, J.M., Franco, D., Domínguez, J., Sineiro, J., Domínguez, H., Núñez, M.J. and Parajó, J.C. 2001. Natural antioxidants from residue sources. *Food Chem.* 72: 145-171.
- Moure, A., Franco, D., Sineiro, J., Domínguez, H., Nunez, M.J. and Lema, J.M. 1999. Antioxidant properties of *Gevuina avellana* hulls. 8th Mediterranean Congress on Chemical Engineering. Barcelona. 10-12.
- Muramatsu, K., Fukuyo, M. and Hara, Y. 1986. Effect of green tea catechins on plasma cholesterol levels in cholesterol-fed rats. *J. Nutr. Sci Vitaminol.* 32: 613-622.
- Nakayama, T., Osawa, T., Mendoza, E.N.T., Laurena, A.C. and Kawakishi, S. 1994. Comparative Study of Antioxidative Assays of Plant Materials. *In Postharvest Biochemistry of Plant Food-Materials in the Tropics.* (Uritani, I., Garcia, V.V. and Mendoza, E.M.T., eds.) p. 241-251. Japan Scientific Societies Press. Japan.
- Namiki, M. 1990. Antioxidants/Antimutagens in foods. *Crit. Rev. Food Sci. Nutr.* 29: 273-300.
- Nawar, W.W. 1996. Lipid. *In Food Chemistry* (Fennema, O.R., ed.). p. 210-243. New York, Marcel Dekker. USA.
- Negre-Salvayre, A. and Salvayre, R. 1992. Quercetin prevents the cytotoxicity of oxidized LDL on lymphoid cell lines. *Free Rad. Biol. Med.* 12: 101.
- Niki, E., Saito, T., Kawakami, A. and Kamiya, Y. 1984. Inhibition of oxidation of methyl linoleate in solution by vitamin E and vitamin C. *J. Biol. Chem.* 259: 4177.

- Nishina, A., Kubota, K., Kameoka, H. and Osawa, T. 1991. Antioxidizing component, musizin, in *Rumex japonicus* Houtt. J. Am. Oil Chem. Soc. 68: 735-739.
- Ohara, Y., Petreson, T.E., Harrison, D.G. 1993. Hypercholesterolemia increases endothelial superoxide anion production. J. Clin. Invest. 91: 2546-2551.
- Olcott, H.S. and Van-Veen. 1963. Role of individual phospholipids as antioxidants. J. Food Sci. 28: 313-315.
- Onyeneho, S.N. and Hettiarachchy, N.S. 1991. Effect of navy bean hull extract on the oxidative stability of soy and sunflower oils. J. Agric. Food Chem. 39:1701-1704.
- Osawa, T. 1994. Novel Natural Antioxidants for Utilization in Food and Biological System. *In* Postharvest Biochemical of Plant Food-Material in the Tropics. (Uritani, I., Garcia, V.V. and Mendoza, E.M.T., eds). p. 241-251. Japan Scientific Societies Press. Japan.
- Osawa, T., Katsuzaki, H., Hagiwara, Y., Hagiwara, H. and Shibamoto, T. 1992. A novel antioxidant isolated from young green barley leaves. J. Agric. Food Chem. 40: 1135-1138.
- Oyaizu, M. 1986. Studies on products of browning reaction: antioxidative activity of products of browning reaction prepared from glucosamine. J. Nutr. 44: 307-315.
- Patsch, W. and Gotto, A.A. 1995. High-Density Lipoprotein Cholesterol, Plasma Triglyceride and Coronary Heart Disease: Pathophysiology and Management. *In* Advances in Pharmacology. (August, J.T., Anders, M.W., Murad, F. and Coyle, J., eds.). p. 375-426. Academic Press. London.
- Pietta, P. and Simonetti, P. 1999. Dietary Flavonoids and Interaction with Physiologic Antioxidant. *In* Antioxidant Food Supplements in Human Health. (Packer, L., Hiramatsu, M. and Yoshikawa, T., eds.). p. 283-308. Academic Press. London.
- Porter, W.L. 1980. Recent trends in food applications of antioxidants. *In* Autoxidation in Food and Biological Systems. (Simic, M.G. and Karel, M., eds.). p. 295-365. Plenum Press. New York.
- Pryzbylski, R., Lee, Y.C. and Eskin, N.A.M. 1998. Antioxidant and radical-scavenging activities of buckwheat seed components. J. Am. Oil Chem. Soc. 75: 1595-1601.

- Puppo, A. 1992. Effect of flavonoids on hydroxyl radical formation by fenton-type reactions; influence of the iron chelator. *Phytochem.* 31: 85-88.
- Rajalakshmi, D. and Narasimhan, S. 1996. Food Antioxidants: Sources and Method of Evaluation. *In Food Antioxidants.* (Madhavi, D.L., Despande, S.S. and Sulunkhe, D.K., eds.), p. 65-158. Marcel Dekker. New York.
- Ramarathnam, N., Osawa, T., Namiki, M. and Kawakishi, S. 1988. Chemical studies on novel rice hull antioxidants. 1. Isolation, fractionation, and partial characterization. *J. Agric. Food Chem.* 36: 719-726.
- Ramarathnam, N., Osawa, T., Namiki, M. and Tashiro, T. 1986. Studies on the relationship between antioxidative activity of rice hull and germination ability of rice seeds. *J. Sci. Food Agric.* 37: 719-726.
- Reio, L. 1958. A method for the paper-chromatographic separation and identification of phenol derivatives, mould metabolism and related compounds of biochemical interest using reference system. *J. Chromatogr.* 1: 338-340.
- Rodriguez de Sotillo, D., Hadley, M. and Holm, E.T. 1994. Phenolics in aqueous potato peel extract, extraction, identification and degradation. *J. Food Sci.* 59: 649-651.
- Saija, A., Scalese, M., Lanza, M., Marzullo, D., Bonina, F. and Castelli, F. 1995. Flavonoids as antioxidant agents: importance of their interaction with biomembranes. *Free Rad. Biol. Med.* 19: 481-486.
- Salas, J.J., Williams, M., Harwood, J.L. and Sanchez. 1999. Lipoxygenase activity in olive (*Olea europaea*) fruit. *J. Am. Oil Chem. Soc.* 76: 1163-1168.
- Sant' Ana, L.S. and Mancini-Filho, J. 2000. Influence of the addition of antioxidants *in vivo* on the fatty acid composition of fish fillets. *Food Chem.* 68: 175-178.
- Shahidi, F. and Alexander, 1998. Green tea cathchins as inhibitors of oxidation of meat lipids. *J. Food Lipid.* 5: 125-133.
- Shahidi, F., Amarovicz. R., Abou-Gharbia, H.A. and Shehata, A.A.Y. 1997. Endogenous antioxidants and stability of sesame oil as affected by processing and storage. *J. Am. Oil Chem. Soc.* 74: 143-147.

- Shahidi, F. and Wanasundara, P.D. 1992. Phenolic antioxidants. *Crit. Rev. Food Sci. Nutr.* 32: 67-103.
- Sherwin, E.R. 1978. Oxidation and antioxidants in fat and oil processing. *J. Am. Oil Chem. Soc.* 55: 809-814.
- Shi, X., Dalal, N.S. and Tain, A.C. 1991. Antioxidant behavior of caffeine: efficient scavenging of hydroxyl radical. *Food Chem.* 29: 1-6.
- Shimada, K., Fujikawa, K., Yahara, K. and Nakamura, T. 1992. Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin. *J. Agric. Food Chem.* 40: 945-948.
- Stahl, W., Junghans, A., de Boer, B., Driomina, E.S., Briviba, K. and Sies, H. 1998. Carotenoid mixtures protect multilamellar liposomes against oxidative damage: synergistic effects of lycopene and lutein. *FEBS Letters.* 427: 305-308.
- Steinberg, D. 1997. Low density lipoprotein oxidation and its pathobiological significance. *J. Biol. Chem.* 272: 20963.
- Taga, M.S., Miller, E.E. and Pratt, D.E. 1984. Chia seeds as source of natural lipid antioxidant. *J. Am. Oil Chem. Soc.* 61: 928-931.
- Terao, J. 1999. Dietary Flavonoids as Plasma Antioxidants on Lipid Peroxidation: Significance of Metabolic Conversion. *In Antioxidant Food Supplements in Human Health.* (Packer, L., Hiramatsu, M. and Yoshikawa, T., eds.). p. 255-268. Academic Press. London.
- Thabrew, M.I., Haghes, R.D. and Mcfarlane, I.C. 1998. Antioxidant activity of *Osbeckia aspera*. *Phytother. Res.* 12: 288-290.
- Tsuda, T., Osawa, T., Nakayama, T., Kawakishi, S. and Ohshima, K. 1993. Antioxidant activity of pea bean (*Phaseolus vulgaris* L.) extract. *J. Am. Oil Chem. Soc.* 70: 909-913.
- Tsuda, T., Ohshima, K., Kawakishi, S. and Osawa, T. 1994a. Antioxidant pigments isolated from the seeds of *Phaseolus vulgaris*. *J. Agri. Food Chem.* 42: 248-251.

- Tsuda, T., Watanabe, M., Ohshima, K., Yamamoto, A., Kawakishi, S. and Osawa, T. 1994b. Antioxidative components isolated from the seed of tamarind (*Tamarindus indica* L.). *J. Agric. Food Chem.* 42: 2671-2674.
- Tsutsumi, K., Kawauchi, Y., Kondo, Y., Inoue, Y., Koshitani, O. and Kohri, Hideaki. 2000. Water extract of defatted rice bran suppresses visceral fat accumulation in rats. *J. Agric. Food Chem.* 48: 1653-1656.
- Ueda, J., Saito, N., Shimazu, Y. and Ozawa, T. 1996. A comparison of scavenging abilities of antioxidants against hydroxyl radicals. *Arch. Biochem. Biophys.* 333: 377-384.
- Uri, N. 1961. Physico-chemical aspects of autoxidation. *In* Autoxidation and Antioxidants. (Lundberg, W.O., ed.). p. 55-106. John Wiley & Sons. New York.
- Ursini, F., Zamburlini, A. and Cazzolato, G. 1998. Postprandial plasma lipid hydroperoxides: a possible link between diet and atherosclerosis. *Free Rad. Biol. Med.* 25: 250.
- Vinson, J.A. and Dabbagh, Y.A. 1998. Effect of green and black tea supplementation on lipids, lipid oxidation and fibrinogen in the hamster: mechanisms for the epidemiological benefits of tea drinking. *FEBS. Lett.* 433: 44.
- Von Schler, A. and Loschner, D. 1985. Natürliche Antioxidantien V, Antioxidantien und Synergisten aus antarktischen Krill. *Fette Seifen Anstrichm.* 87: 454-457.
- Wanasundara, U.N. and Shahidi, F. 1998. Antioxidant and pro-oxidant activity of green tea extracts in marine oils. *Food Chem.* 63: 335-342.
- Wang, S.Y. and Lin, H.S. 2000. Antioxidant activity in fruits and leaves of blackberry, raspberry, and strawberry varies with cultivar and developmental stage. *J. Agric. Food Chem.* 48: 140-146.
- Wettasingha, M. and Shahidi, F. 1999. Antioxidant and free radical scavenging properties of ethanolic extracts of defatted borage (*Borago officinalis* L.) seeds. *Food Chem.* 67: 399-414.
- Weurman, C. and Swain, T. 1955. Changes in the enzymatic browning of barley's seedling applied during their development. *J. Sci. Food Agric.* 6: 186-190.

- Whalley, C.V., Rankin, S.M., Houlst, J.R.S., Jessup, W. and Leake, D.S. 1990. Flavonoids inhibit the oxidative modification of low density lipoproteins by macrophages. *Biochem. Pharmacol.* 39: 1743.
- Williams, P., Ongsakul, M., Proudfoot, J., Croft, K. and Beilin, L. 1995. Mangostin inhibits the oxidative modification of human low density lipoprotein. *Free Rad. Res.* 23: 175-184.
- Willson, K. 1999. Botany and Plant Improvement. *In Crop Production Science in Horticulture.* (Atherton, J., ed.). p. 167-173. Cabi. New York.
- Wiseman, S., Weisgerber, U., Tijburg, L. and Korver, O. 1999. The Food Industry and Functional Foods: Tea Antioxidants and Cardiovascular Disease. *In Antioxidant Food Supplements in Human Health.* (Packer, L., Hiramatsu, M. and Yoshikawa, T., eds.). p. 445-460. Academic Press. London.
- Yamaguchi, F., Yoshimura, Y., Nakazawa, H. and Ariga, T. 1999. Free radical-scavenging activity of grape seed extract and antioxidants by electron spin resonance spectrometry in an $H_2O_2/NaOH/DMSO$ system. *J. Agric. Food Chem.* 47: 2544-2548.
- Yamaguchi, T., Takamura, H., Matoba, T. and Terao, J. 1998. HPLC method for evaluation of the free radical-scavenging activity of foods by using 1,1-diphenyl-2-picrylhydrazyl. *Biosci. Biotechnol. Biochem.* 62: 1201-1204.
- Yang, J.H., Mau, J.L., Ko, P.T. and Huang, L.C. 2000. Antioxidant properties of fermented soybean broth. *Food Chem.* 71: 249-254.
- Yang, T.T.C and Koo, M.W.L. 2000. Inhibitory effect of Chinese green tea on endothelial cell-induced LDL oxidation. *Atherosclerosis.* 148: 67-73.
- Yen, G.C. and Chen, H.Y. 1995. Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J. Agric. Food Chem.* 43: 27-32.
- Yen, G.C. and Chuang, D.Y. 2000. Antioxidant properties of water extracts from *Cassia tora* L. in relation to the degree of roasting. *J. Agric. Food Chem.* 48: 2760-2765.
- Yen, G.C. and Duh, P.D. 1993. Antioxidative properties of methanolic extracts from peanut hulls. *J. Am. Oil Chem. Soc.* 70: 383-386.

- Yen, G.C. and Duh, P.D. 1994. Scavenging effect of methanolic extracts of peanut hulls on free-radical and active-oxygen species. *J. Agric. Food Chem.* 42: 629-632.
- Yen, G.C. and Lee, C.E. 1997. Antioxidative properties of extracts from *Aspergillus candidus* broth filtrate. *J. Sci. Food Agric.* 75: 326-332.
- Yen, G.C. and Wu, J.Y. 1999. Antioxidant and radical-scavenging properties of extracts from *Ganoderma tsugae*. *Food Chemistry.* 65: 375-379.
- Yen, G.C., Chen, H.Y. and Peng, H.H. 1997. Antioxidant and Pro-oxidant effects of various tea extracts. *J. Agric. Food Chem.* 45: 30-34.
- Yen, G.C., Duh, P.D. and Chuang, D.Y. 2000. Antioxidant activity of anthraquinones and anthrone. *Food Chem.* 70: 437-441.
- Yen, G.C., Wu, S.C. and Duh, P.D. 1996. Extraction and identification of antioxidant component from the leaves of mulberry (*Morus alba* L.). *J. Agric. Food Chem.* 44: 1687-1690.
- Yi, O.S., Han, D. and Shin, H.K. 1991. Synergistic antioxidative affects of tocopherol and ascorbic acid in fish oil/lecithin/water system. *J. Am. Oil Chem. Soc.* 68: 881-883.
- Yoshida, H., Ishikawa, T., Hosoi, H. 1999. Inhibitory effect of tea flavonoids on the ability of cells to oxidize low density lipoprotein. *Biochem. Pharmacol.* 58 : 1695.
- Yoshioka, H., Sugiura, K., Kawahara, R., Fujita, T., Makino, M., Kamiya, M. and Tsuyumu, S. 1991. Formation of radicals and chemiluminescence during the autoxidation of tea catechins. *Agric. Biol. Chem.* 55: 2717-2723.
- Zarzuelo, A. 1991. Vasodilator effect of olive leaf. *Planta Medica.* 57: 417-419.
- Zhang, A., Chan, P.T., Luk, Y.S., Ho, W.K.K and Chen, Z.Y. 1997. Inhibitory effect of jasmine green tea epicatechin isomers on LDL-oxidation. *J. Nutr. Biochem.* 8: 334-340.
- Zheng, T.Z., Tan, Y.F., Huang, G.X., Fan, H. and Ma, B. 1988. Mulberry cultivation. *FAO Agricultural Services Bull.* 73/1: 1-20.

- Zhishen, J., Mengcheng, T. and Jianming, W. 1999. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem.* 64: 555-559.
- Zhu, N., Wang, M., Wei, G.J., Lin, J.K., Yang, C.S. and Ho, C.T. 2000a. Identification of reaction products of (-)-epigallocatechin, (-)- epigallocatechin gallate and pyrogallol with 2,2-diphenyl-1-picrylhydrazyl radical. *Food Chem.* 73: 345-349.
- Zhu, Q.Y., Huang, Y. and Chen, Z.Y. 2000b. Interaction between flavonoids and α -tocopherol in human low density lipoprotein. *J. Nutr. Biochem.* 11: 14.
- Zhu, Q.Y., Zhang, A., Tsang, D., Huang, Y. and Chen, Z.Y. 1997. Stability of green tea catechins. *J. Agric. Food Chem.* 45: 4624-4628.