

REFERENCES

- Ahn, D.H., Hattori, A. and Takahashi, K. 1993. Structural changes in Z-disks of skeletal muscle myofibrils during growth of chicken. *J. Biochem.* 113: 383-388.
- Ajuyah, A.O., Hardin, R.T., Cheung, K. and Sim, J.S. 1992. Yield, lipid, cholesterol and fatty acid composition of spent hens fed full-fat oil seeds and fish meal diets. *J. Food Sci.* 57: 338-341.
- Allen, C.D., Fletcher, D.L., Northcutt, J.K. and Russel, S.M. 1998. The relationship of broiler breast color to meat quality and shelf-life. *Poultry Sci.* 77: 361-366.
- AOAC. 1999. Official Methods of Analysis. 16th Ed. Association of Official Analytical Chemists, Washington, DC.
- Argonasa, G.C. and Marriott, N.G. 1989. Organic acids as tenderizers of collagen in restructured beef. *J. Food Sci.* 54: 1173-1176.
- Bailey, A.J. 1972. The basis of meat texture. *J. Sci. Food Agric.* 23: 995-1007.
- Bailey, A.J. and Light, N.D. 1989. *Connective Tissue in Meat and Meat Products*. Elsevier Applied Science. New York.
- Bendal, J.R. 1979. Relations between muscle pH and important biochemical parameters during the post-mortem changes in mammalian muscles. *Meat Sci.* 3: 143-157.
- Berge, P., Ertbjerg, P., Larsen, L.M., Astruc, T., Vignon, X. and Moller, A.J. 2001. Tenderization of beef by lactic acid injected at different times post mortem. *Meat Sci.* 57: 347-357.
- Bergman, I. and Loxley, R. 1963. Two improved and simplified methods for the spectrophotometric determination of hydroxyproline. *Anal. Chem.* 35: 1961-1965.
- Bertola, N.C., Bevilacqua, A.E. and Zaritzky, N.E. 1994. Heat treatment effect on texture changes and thermal denaturation of proteins in beef muscle. *J. Food Process. Preserv.* 18: 31-46.
- Bircan, C. and Barringer, S.A. 2002. Determination of protein denaturation of muscle foods using the dielectric properties. *J. Food Sci.* 67: 202-205.
- Bouton, P.E., Harri, P.P. and Ratcliff, D. 1981. Effect of cooking temperature and time on the shear properties of meat. *J. Food Sci.* 46: 1082-1086.
- Brody, A. 2003. Food canning in the 21st century. *Food Technol.* 56: 75-79.

- Califano, A.N., Bertola, N.C., Bevilacqua, A.E. and Zaritzky, N.E. 1997. Effect of processing conditions on the hardness of cooked beef. *J. Food Engin.* 34: 41-54.
- Chotsangkad, R. and Kongrattananun, N. 1999. Growth and carcass quality of native chickens raised under the natural day, length and the photoperiod of twenty-three hours a day. *The Kasetsart J.* 33: 60-74. (In Thai)
- Christensen, M., Purslow, P.P. and Lersen, L.M. 2000. The effect of cooking temperature on mechanical properties of whole meat, single muscle fibres and perimysial connective tissue. *Meat Sci.* 55: 301-307.
- Chrystall, B. 1994. Meat texture measurement. *In Quality Attributes and their Measurement in Meat, Poultry and Fish Products* (Pearson, A.M. and Dutson, T.R., eds.). p. 316-336. Black Academic & Professional. UK.
- Conforth, D. 1994. Color its basis and importance. *In Quality Attributes and their Measurement in Meat, Poultry and Fish Products* (Pearson, A.M. and Dutson, T.R., eds.). p. 34-78. Black Academic & Professional. UK.
- Dawson, P.L. 1997. High temperature processing effects on the properties of fowl. *Poultry Sci.* 76: 774 -779.
- Dawson, P.L., Sheldon, B.W. and Miles, J.J. 1991. Effect of aseptic processing on the texture of chicken meat. *Poultry Sci.* 70: 2359-2367.
- Department of Livestock Development. 2003. Data of Livestock Economic for the Year 2003. Bangkok. (In Thai)
- Department of Livestock Development. 2007. Data of Livestock Economic for the Year 2007. Bangkok. (In Thai)
- Deumier, F. 2004. Pulsed-vacuum immersion of chicken meat and skin in acid solutions. Effect on mass transfers, colour and microbial quality. *Int. J. Food Sci. Tech.* 39: 277-286.
- Ding, H., Xu, R.J. and Chan, D.K.O. 1999. Identification of broiler chicken meat using a visible/near-infrared spectroscopic technique. *J. Sci. Food Agric.* 79: 1382-1388.
- Doerscher, D.R., Briggs, J.L. and Longernan, S.M. 2003. Effect of pork collagen on the thermal and viscoelastic properties of purified myofibrillar protein gels. *J. Food Sci.* 66: 181-188.

- Dransfield, E. 1994. Tenderness of meat, poultry and fish. *In* Quality Attributes and their Measurement in Meat, Poultry and Fish Products (Pearson, A.M. and Dutson, T.R., eds.). p. 289-315. Black Academic & Professional. UK.
- Duun, A.A., Kilpatrick, D.J. and Gault, N.F.S. 1993. Effect of post-mortem temperature on chicken M. pectoralis major muscle shortening and cooked meat tenderness. *Br. Poultry Sci.* 34: 689-697.
- Dunn, A.A., Tolland, E.L.C., Kilpatrick, D.J. and Gault, N.F.S. 2000. Relationship between early post-mortem muscle pH and shortening induced toughness in the Pectoralis major muscle of processed broilers air chilled at 0°C and 12°C. *Br. Poultry Sci.* 41: 53-60.
- Eilert, S.J. and Mandigo, R.W. 1993. Procedure for soluble collagen in thermally processed meat products. *J. Food Sci.* 58: 948-949.
- El, S.N. 1995. Evaluating protein quality of meats using collagen content. *Food Chem.* 53: 209-210.
- Fernandez, X., Monin, G., Talmant, A., Mourot, J. and Lobret, B. 1999. Influence of intramuscular fat content on the quality of pig meat. Composition of the lipid fraction and sensory characteristic of *m. Longissimus lumborum*. *Meat Sci.* 53: 59-65.
- Fletcher, D.L. 1999a. Poultry meat color. *In* Poultry Meat Science, Poultry Science Symposium Series (Richardson, R.I. and Mead, G.C., eds.). p. 159-175. CABI Publishing. UK.
- Fletcher, D.L. 1999b. Broiler breast meat color variation, pH and texture. *Poultry Sci.* 78: 1323-1327.
- Fletcher, D.L., Qiao, M. and Smith, D.P. 2000. The relationship of raw broiler breast meat color and pH to cooked meat color and pH. *Poultry Sci.* 79: 784-788.
- Foegeding, E.A., Lanier, T.C. and Hultin, H.O. 1996. Characteristics of edible muscle tissues. *In* Food Chemistry (Fennema, O.R., ed.). p. 902-906. Marcel Dekker, Inc. New York.
- Gault, N.F.S. 1984. The influence of acetic acid concentration on the efficiency of marinading as a process for tenderizing beef. *In* Meat Research Workers^{3rd} European Meeting. Bristol.
- Gault, N.F.S. 1985. The relationship between water-holding capacity and cooked meat tenderness in some beef muscles as influenced by acidic conditions below the ultimate pH. *Meat Sci.* 15: 15-30.

- Gault, N.F.S. 1991. Marinated meat. *In* Development in meat science (Lawrie, R.A. ed.). p. 191-246. Elsevier Applied Science. UK.
- Goll, D.E., Bray, R.W. and Hoekstra, W.G. 1963. Age associated changes in muscle composition. *J. Food Sci.* 28: 503-509.
- Hamm, R. 1960. Biochemistry of meat hydration. *In* Advances in Food Research (Chichester, C.O., Mrak, E.M. and Stewart, G.F., eds.). p. 355-463. Academic Press. New York.
- Hamm, R. 1977. Changes in muscle proteins during the heating of meat. *In* Physical, Chemical and Biological Changes in Food Caused by Thermal Processing, (Hoyem, T. and Kvale, O., eds.). p. 101-134. Applied Science Publisher. London.
- Heerden, S.M., Schönfeldt, H.C., Smith, M.F. and Jansen van Rensburg, D.M. 2002. Nutrient content of South African Chickens. *J. Food Comp Anal.* 15: 47-64.
- Honikel, K.O. and Hamm, R. 1994. Measurement of water-holding capacity and juiciness in important Mediterranean finfish, crustaceans and mollusks. *J. Food Comp Anal.* 8: 258-273.
- Karakoltsidis, P.A., Zaotos, A. and Constantinides, S.M. 1998. Composition of the commercially important Mediterranean finfish, crustaceans and mollusks. *J. Food Compo. Anal.* 8: 258-273.
- Kijowski, J.M. and Mast, M.G. 1988. Thermal properties of proteins in chicken broiler tissue. *J. Food Sci.* 53: 363-366.
- Kivikari, R. 1996. Buffering capacity of meat. Ph.D. Thesis. University of Helsinki, Finland.
- Koohmaraie, M., Whipple, G., Kretchmar, D.H., Crouse, J.D. and Mersmann, H.J. 1992. Postmortem proteolysis in longissimus muscle from beef, lamb and pork carcasses. *J. Animal Sci.* 69: 617-624.
- Kyla-Puhju, M., Ruusunen, M., Kivikari, R. and Puolanne, E. 2004. The buffering capacity of porcine muscles. *Meat Sci.* 67: 587-593.
- Lawrie, R. A. 1991. Meat Science. Oxford Pergamon Press.
- Lee, S.O., Min, J.S. and Lee, M. 2003. Physical evaluation of popped cereal snacks with spent hen meat. *Meat Sci.* 64: 383-390.
- Li, C.T. 2006. Myofibrillar protein extracts from spent hen meat to improve whole muscle processed meat. *Meat Sci.* 72: 581-583.

- Liu, A., Nishimura, T. and Takahashi, K. 1994. Structural changes in endomysium and perimysium during post-mortem ageing of chicken semitendinosus muscle contribution of structural weakening of intramuscular connective tissue to meat tenderization. *Meat Sci.* 38: 315-328.
- Liu, H.J., Nishimura, T. and Takahashi, K. 1996. Relationship between structural properties of intramuscular connective tissue and toughness of various chicken skeletal. *Meat Sci.* 43: 43-49.
- Maran, E.T. 1999. Live production factors influence yield and quality of poultry meat. *In Poultry Meat Sci.* p.179-195. Oxon CABI Publishing.
- Marten, H., Stabursvik, E. and Marten, M. 1982. Texture and colour changes in meat during cooking related to thermal denaturation of muscle proteins. *J. Texture Stud.* 13: 291-309.
- McCarthy, M.J. Heli, J.R., Kruegermann, C. and Desvignes, D. 1991. Acid requirement for pH modification of processed foods. *J. Food Sci.* 56: 973-976.
- Mehaffey, J.M., Prandhan, S.P., Meullenet, J.F., Emmert, L., McKee, S.R. and Owen, C.M. 2006. Meat quality evaluation of minimally aged broiler breast fillets from five commercial genetic strains. *Poultry Sci.* 85: 902-908.
- Meyer, L.H. 1973. Meat and meat product. *In Food Chemistry* (Sisler, H.S. and Vanderwereds, C.A., eds.). Press Pvt. LTD. New Delhi.
- Miles, C.L. and Lawrie, R.A. 1970. Relation between pH and tenderness in cooked muscle. *Int. J. Food. Sci.* 54: 325-330.
- Miller, R.K. 1994. Quality characteristic. *In Muscle Foods* (Kinsman, D.M., Kotula, A.W. and Breidenstein, B.C., eds.). Chapman & Hall. New York.
- Moreira, L.A., Oliveira, F.A.R. and Silva, T.R. 1992. Prediction of pH change in processed acidified turnips. *J. Food Sci.* 57: 928-931.
- Murphy, R.Y. and Marks, B.P. 2000. Effect of meat temperature on properties, texture and cook loss for ground chicken breast patties. *Poultry Sci.* 79: 99-104.
- Murphy, R.Y., Marks, B.P. and Marcy, J.A. 1998. Apparent specific heat of chicken breast patties and their constituent proteins by differential scanning calorimetry. *J. Food Sci.* 63: 88-91.
- Nakamura, R., Sekoguchi, S. and Sato, Y. 1975. The contribution of intramuscular collagen to the tenderness of meat from chickens with different ages. *Poultry Sci.* 54: 1604-1612.

- Naveena, B.M. and Mendiratta, S.V. 2001. Tenderisation of spent hen meat using ginger extract. *Br. Poultry Sci.* 42: 344-349.
- Nishida, J. and Nishida, T. 1985. Relationship between the concentration of myoglobin and parvalbumin in various types of muscle tissues from chickens. *Br. Poultry Sci.* 26: 105-115.
- Noppawan, C., Pojan, A. and Vanasitchaiwat, W. 1998. Effect of protein level and raising condition on growth rate and carcass quality of crossbreed indigenous chicken. *Swine Newsletter.* 23: 55-70. (In Thai)
- Northcutt, J.K. 1997. Factor affecting poultry meat quality. Bulletin 1157: Department of poultry Science, The University of Georgia, Athens, GA.
- Northcutt, J.K., Pringle, T.D., Young, L.L., Buhr, R.J. and Young, L.L. 1998. Effects of age and tissue type on the calpain proteolytic system in turkey skeletal muscle. *Poultry Sci.* 77: 367-372.
- Nowsad, A.A.K.M., Kanoh, S. and Niwa, E. 2000. Thermal gelation characteristics of breast and thigh muscles of spent hen and broiler and their surimi. *Meat Sci.* 54: 169-175.
- Offer, G. and Knight, P. 1988. The structural basis of water-holding capacity in meat. *In Development in Meat Science.* p. 173-243. Elsevier Science Publication. London.
- Offer, G. and Trinick, J. 1983. On the mechanism of water holding in meat the swelling and shrinking of myofibrils. *Meat Sci.* 8: 245-281.
- Okkazaki, E., Kanna, K. and Suzuki, T. 1986. Effect of sarcoplasmic protein on rheological properties of fish meat gel formed by retort-heating. *Bull. Jap. Soc. Fish.* 52: 1821-1827.
- Ozawa, S., Mitsunashi, T., Mitsumoto, M., Matsumoto, S., Itoh, N., Itagaki, K., Kohno, Y. and Dohgo, T. 2000. The characteristics of muscle fiber types of *Longissimus thoracis* muscle and their influences on the quantity and quality of meat from Japanese black steer. *Meat Sci.* 54: 65-70.
- Palka, K. and Daun, H. 1999. Changes in texture, cooking losses, and myofibrillar structure of bovine *M. semitendinosus* during heating. *Meat Sci.* 51: 237-243.
- Palka, K. 1999. Changes in intramuscular connective tissue and collagen solubility of bovine *M. semitendinosus* during retorting. *Meat Sci.* 53: 189-194.

- Pearson, A.M. and Young, R.B. 1989. Muscle and Meat Biochemistry. Academic Press, Inc. California.
- Penny, I.F., Voyle, C.A. and Lawrie, R.A. 1963. A comparison of freeze-dried beef muscles of high or low ultimate pH. *J. Sci. Food Agric.* 14: 535-545.
- Plowman, J.E. and Close, E.A. 1988. An evaluation of a method to differentiate the species of origin of meats on the basis of the contents anserine, balenine and carnosine in skeletal muscle. *J. Sci. Food Agri.* 45: 69-78.
- [Rao](#), M.A. and Lund, D.B. 1986. Kinetics of thermal softening of foods. *J. Food Process. Pres.* 10: 311-329.
- Rao, M.V., Gault, N.F. and Kennedy, S. 1989. Changes in the ultrastructure of beef muscles as influenced by acidic conditions below the ultimate pH. *Food Microstruct.* 8: 115-124.
- Rochidi, A., Foucat, L. and Renou, J. 2000. NMR and DSC studies during thermal denaturation of collagen. *Food Chem.* 69: 295-299.
- Sams, R. 1990. Lathrogen effects on the collagen heat stability and tenderness of spent fowl muscles. *Poultry Sci.* 69: 477-481.
- Sanchai, J., Leangwunta, V., Leotaragul, A. and Worachai, L. 2003. A comparative study of Thai native chicken and broiler on productive performance, carcass and meat quality. Research Report Chiang Mai University. (In Thai)
- Saunders, A.B. 1994. The effect of acidification on myofibrillar proteins. *Meat Sci.* 37: 271-280.
- Shorthose, W.R. and Harris, P.V. 1990. Effect of animal age on the tenderness of selected beef muscles. *J. Food Sci.* 55: 253-261.
- Sims, T.J. and Bailey, A.J. 1981. Connective tissue. *In* Developments in Meat Science (Lawrie, R.A., ed.), p. 29-59. Applied Science Publishers.
- Singh, R.R.B., Rao, K.H., Anjaneyulu, A.S.R. and Patil, G.R. 2001. Moisture sorption properties of smoked chicken sausages from spent hen meat. *Food Res. Inter.* 34: 143-148.
- Siripongvutikorn, S. 2004. Postharvest deterioration of Seabass (*Lates calcarifer*) and its shelf-life extension by using Tom-Yum mix. Ph.D. Dissertation. Prince of Songkla University.
- Siripongvutikorn, S., Thummaratwasik, P. and Huang, Y. 2005. Antimicrobial and antioxidant effect of Thai seasoning, Tom Yum. *LWT.* 38: 347-352.

- Smith, D.P. and Fletcher, D.L. 1988. Chicken breast muscle fiber type and diameter as influenced by age and intramuscular location. *Poultry Sci.* 67: 908-913.
- Smith, D.P., Fletcher, D.L., Buhr, R.J. and Beyer, R.S. 1993. Perkin duckling and broiler chicken *pectoralis* muscle structure and composition. *Poultry Sci.* 72: 202-208.
- Stabursvik, E. and Martens, H. 1980. Thermal denaturation of proteins in post rigor muscle tissue as studied by differential scanning calorimetry. *J. Sci. Food Agric.* 31: 1034-1038.
- Stell, R.G.D. and Torrie, J.H. 1980. Principles and procedures of statistics: A Biometrical Approach. 2nd ed. New York: McGraw-Hill.
- Stewart, G.F. and Amarine, M.A. 1982. Introduction to Food Science and Tecnology. 2nd ed. Academic Press. New York.
- Svensson, A.B. and Tonberg, E. 1998. Prediction of acid induced pH decrease in meat. *In* Proceeding of the 44th ICoMST Barcelona Spain. p. 506-507.
- Swatland, H. J. 1984. Structure and Development of Meat Animals. Prentice-Hall Inc.
- Torrescano, G., Sanchez-Escalante, A., Gimenez, B., Roncales, P. and Beltran, J.A. 2003. Shear values of raw samples of 14 bovine muscles and their relation to muscle collagen characteristics. *Meat Sci.* 64: 85-91.
- Twiddy, D.R., Cross, S.J. and Cooke, R.D. 1987. Parameters involved in the production of lactic acid preserved fish-starchy substrate combinations. *Int. J. Food Sci. Tech.* 22: 115-121.
- Van Marle-Koster, E. and Webb, E.C. 2000. Carcass characteristics of South African native chicken lines. *South African J. Animal Sci.* 30: 53-56.
- Voller, L.M., Dawson, P.L. and Han, I.Y. 1996. Processing temperature and moisture content effects on the texture and microscopic appearance of cooked fowl meat gels. *Poultry Sci.* 75: 1603-1610.
- Voller-Reasonover, L., Han, I.Y., Acton, J.C., Titus, T.C., Bridges, W.C. and Dawson, P.L. 1997. High temperature processing effects on the properties of fowl meat gels. *Poultry Sci.* 76: 774-779.
- Warriss, P.D. 2000. Meat Science: An introductory Text. Oxon: CABI.
- Wattanachant, S. and Wattanachant, C. 2007. Chemical composition, properties and microstructure of Thai indigenous chicken muscle as influenced by age rearing systems. Research Report. Prince of Songkla University.

- Wattanachant, S., Benjakul, S. and Ledward, D.A. 2004. Compositions, color and texture of Thai indigenous and broiler chicken muscles. *Poultry Sci.* 83: 123-128.
- Wattanachant, S., Benjakul, S. and Ledward, D.A. 2005a. Microstructure of Thai indigenous and broiler chicken muscles. *Poultry Sci.* 84: 328-326.
- Wattanachant, S., Benjakul, S. and Ledward, D.A. 2005b. Effect of heat treatment on changes in texture structure and properties of Thai indigenous chicken muscles. *Food Chem.* 93: 337-348.
- Wu, F.Y., Tsao, P.H., Lee, Y.C. and Guo, S.F. 1996. Effect of acid or alkaline treatment on thermal denaturation of connective tissue from spent hen. *Chinese Animal Sci.* 25: 335-345.
- Xiong, S., Xiong, Y.L., Blanchard, S.P., Wand, B. and Tidwell, J.H. 2002. Evaluation of tenderness in prawns (*Machrobranchium rosenbergii*) marinated in various salt and acid solution. *Int. J. Food Sci. Technol.* 37: 291-296.
- Xiong, Y.L. and Brekke, C.J. 1989. Thermal denaturation of muscle proteins from different species and muscle types as studied by differential scanning calorimetry. *Can. Inst. Food Sci. Technol. J.* 20: 357-362.
- Xiong, Y.L., Cantor, A.H., Pescatore, A.J., Blanchard, S.P. and Straw, M.L. 1993. Variations in muscle chemical composition pH and protein extractability among eight different broiler crosses. *Poultry Sci.* 72: 583-588.
- Xiong, Y.L., Ho, C.T. and Shahidi, F. 1999. Quality characteristics of muscle foods. *In* Quality Attributes of Muscle Foods (Xiong, Y.L., Ho, C.T. and Shahidi, F., eds.). Kluwer Academic/Plenum Publishers. New York.