

## เอกสารอ้างอิง

- กรมปะรัง. 2541. คู่มือการเพาะเลี้ยงปลานิลเพศผู้ชายพันธุ์จีตราด 2. กรุงเทพฯ: สถาบันวิจัยและพัฒนาพันธุกรรมสัตว์น้ำ. กรมปะรัง กระทรวงเกษตรและสหกรณ์.
- กรมปะรัง. 2545. สถิติการปะรังแห่งประเทศไทย พ.ศ. 2542. เอกสารฉบับที่ 10/2545. กรุงเทพฯ: กองเศรษฐกิจการปะรัง. กรมปะรัง กระทรวงเกษตรและสหกรณ์.
- กรมปะรัง. 2548. สถิติผลผลิตการเลี้ยงสัตว์น้ำจีดประจำปี 2546. เอกสารฉบับที่ 6/2548. กรุงเทพฯ: กลุ่มวิจัยและวิเคราะห์สถิติการปะรัง. กรมปะรัง กระทรวงเกษตรและสหกรณ์.
- นิรุทธิ์ สุขเกษม. 2544. ผลของระดับการเนื้อเม็ดในปาล์มน้ำมันต่อการเจริญเติบโตของปลานิล (*Oreochromis niloticus* Linn.). วิทยานิพนธ์วิทยาศาสตร์มหาบัณฑิตมหาวิทยาลัยสงขลานครินทร์.
- พรณศรี จริโนกาส. 2531. ปลานิลสีแดงพันธุ์ไทย. ว. การปะรัง 41: 41-43.
- มะลิ บุณยรัตน์ ผลิน และ จุยะดี พงศ์มณีรัตน์. 2533. ความต้องการฟอสฟอรัสในอาหารปลากระเพรา. เอกสารวิชาการ. สงขลา: สถาบันวิจัยการเพาะเลี้ยงสัตว์ชายฝั่ง. กรมปะรัง กระทรวงเกษตรและสหกรณ์.
- มานพ ตั้งตรงไฟโกรจน์, ภาณุ เทวัตโนมณีกุล, พรณศรี จริโนกาส, สุจินต์ หนูขาวัญ, กำชัย ลาภัณยุฑิ, วีระ วัชรกรโยธิน และ วิมล จันทร์โรทัย. 2536. การพัฒนาการเพาะเลี้ยงปลานิล. เอกสารเผยแพร่ฉบับที่ 23. กรุงเทพฯ: สถาบันวิจัยการเพาะเลี้ยงสัตว์น้ำจีด. กรมปะรัง กระทรวงเกษตรและสหกรณ์.
- วีรพงศ์ วุฒิพันธุ์ชัย. 2536. การเพาะพันธุ์ปลา. ภาควิชาวิชาศาสตร์ คณะวิทยาศาสตร์ มหาวิทยาลัยปูรพา.
- วุฒิพrho พรมขุนทอง และ อัจฉริยา มูสโกกาส. 2548. ผลของเอนไซม์ไฟเตสต์ต่อการเพิ่มการใช้วัตถุดิบพืชในปลานิลแดงแบล็งเพส (*Oreochromis niloticus* Linn.) ว. สงขลานครินทร์ วทท. (ฉบับพิเศษ 1) 27: 151-170.
- วุฒิพrho พรมขุนทอง, วรรณชัย พรมเกิด, กิจการ ศุภมาตย์, วุฒิกรรณ์ จิตติวรรณ และ ดุสิต นาคะชาต. 2547. การแทนที่ปลาป่นในอาหารปลานิลแดงแบล็งเพส (*Oreochromis niloticus* Linn.) ด้วยการกneื้อเม็ดในปาล์มน้ำมัน. ว. สงขลานครินทร์ วทท. 26: 167-179.

สำนักงานคณะกรรมการพัฒนาการเศรษฐกิจและสังคมแห่งชาติ. 2544. แผนพัฒนาเศรษฐกิจและสังคมฉบับที่ 9 พ.ศ. 2545-2549. เข้าถึงได้จาก [http://www.Mesdb.go.th/plam/data/plan9/contemt\\_page.html](http://www.Mesdb.go.th/plam/data/plan9/contemt_page.html). เข้าถึงเมื่อ 18 มกราคม 2548.  
อุทัยรัตน์ ณ นคร. 2539. การเพาะขยายพันธุ์ปลา. ภาควิชาเพาะเลี้ยงสัตว์น้ำ. กรุงเทพฯ: คณะประมง มหาวิทยาลัยเกษตรศาสตร์.

- Adeola, O. and Sands, J.S. 2003. Does supplemental dietary microbial phytase improve amino acid utilization? A perspective that it does not. *J. Anim. Sci.* 81: 78–85.
- Alceste, C.C. 2004. Alternative protein sources In tilapia feed formulation. *Aquacult. Mag.* 30: 42-45.
- Amirkolaie, A. K., Leenhouters, J.L., Verreth, J. A. J. and Schrama, J.W. 2005. Type of dietary fibre (soluble versus insoluble) influences digestion, faeces characteristics and faecal waste production in Nile tilapia (*Oreochromis niloticus* L.). *Aquaculf. Res.* 36: 1157-1166.
- Andrews, J.W., Muri, T. and Campbell, C. 1973. Effects of dietary calcium and phosphorus on growth, food conversion, bone ash and hematocrit levels of catfish. *Ictalurus punctatus*. *J. Nutr.* 103: 766-771.
- Austreng, E., Storebakken, T., Thomassen, M.S., Refstie, S. and Thomassen, Y. 2000. Evaluation of selected trivalent metal oxides as inert markers used to estimate apparent digestibility in salmonids. *Aquaculture* 188: 65–78.
- AOAC 1990. Official Methods of Analysis 15<sup>th</sup>. Washington: The Association of official Analytical Chemists.
- Azevedo, P.A., Leeson, S., Cho, CY. and Bureau, D.P. 2004. Growth, nitrogen and energy utilization of juveniles from four salmonid species: diet, species and size effects. *Aquaculture* 234: 393-414.
- Baden, S.P., Loo, L.O., Pihl, L. and Rosenberg, R. 1990. Effects of eutrophication on benthic communities including fish. *Swedish West Coast. AMBIO.* 19: 113-122.

- Baruah, K., Pal, A.K., Sahu, N.P., Jain, K.K., Mukherjee, S.C. and Debnath, D. 2005. Dietary protein level, microbial phytase, citric acid and their interactions on bone mineralization of *Labeo rohita* (Hamilton) juveniles. Aquaculf. Res. 36: 803 - 812.
- Blaxhall, P.C. and Daisley, K.W. 1973. Routine hematological methods for use with fish blood. J. Fish Biol. 5: 771-781.
- Boonyaratpalin, M. and Phromkunthong, W. 2000. Effects of Ronozyme VP treated rice bran and oil palm meal on growth of sex reversed *Tilapia nilotica*. The 6<sup>th</sup> Roche Aquaculture Conference Asia Pacific, Bangkok, Thailand, September 29.
- Borgeson, T.L., Racz, . V.J., Wilkie, D.C., White, L.J. and Drew, M.D. 2006. Effect of replacing fishmeal and oil with simple or complex mixtures of vegetable ingredients in diets fed to Nile tilapia (*Oreochromis niloticus*). Aquacul. Nutr. 12: 141–149.
- Bureau, D.P. 2004. Factors affecting metabolic waste outputs in fish. Avances en Nutrición Acuícola VII Memorías del Septimo Simposium Internacional de Nutrición Acuícola. Centro de las Artes, Universidad de Sonora Hermosillo, Sonora, Mexico, 16-18 Noviembre, 2004.
- Bureau, D.P. and Cho, C.Y. 1999. Phosphorus utilization by rainbow trout (*Oncorhynchus mykiss*): estimation of dissolved phosphorus waste output. Aquaculture 179: 127–140.
- Bureau, D.P., Kirkland, J.B. and Cho, C.Y. 1998. The partitioning of energy from digestible carbohydrate by rainbow trout (*Oncorhynchus mykiss*) In Energy Metabolism of Farm Animals feed (eds K.J. McCracken, E.F. Unsworth and A.R.G. Wylle). Wallingford. UK: CAB International Press.
- Boyd, C.E. and Tucker, C.S. 1992. Water Quality and Pond Soil Analysis for Aquaculture. Alabama: Auburn University.

- Cheng, Z.J. and Hardy, R.W. 2002. Apparent digestibility coefficients of nutrients and nutritional value of poultry by-product meals for rainbow trout *Oncorhynchus mykiss* measured in vivo using settlement. J. World Aquacult. Soc. 33: 458-465.
- Cheng, Z.J. and Hardy, R.W. 2003. Effect of extrusion and expelling processing, and microbial phytase supplementation on apparent digestibility coefficients of nutrients in full-fat soybeans for rainbow trout (*Oncorhynchus mykiss*). Aquaculture 218: 501-514.
- Cheng, Z.J., Hardy, R.W., Verlhac, V. and Gabaudan, J. 2004. Effects of microbial phytase supplementation and dosage on apparent digestibility coefficients of nutrients and dry matter in soybean product-based diets for rainbow trout *Oncorhynchus mykiss*. J. World Aquacult. Soc. 35: 1-15.
- Cho, C.Y., Hynes, J.D., Wood, K.R. and Yoshida, H.K. 1994. Development of high-nutrient-dense, low-pollution diets and prediction of aquaculture wastes using biological approaches. Aquaculture 124: 293-305
- Cho, C.Y. and Bureau, D.P. 2001. A review of diet formulation strategies and feeding systems to reduce excretory and feed wastes in aquaculture. Aquaculf. Res. 32: 349-360.
- Cho, C.Y., Cowey, C.B. and Watanabe, T. 1985. Finfish nutrition in Asia. Methodological approaches to research and development. Ottawa: International Development Research Centre.
- Cho, C.Y., Slinger, S.J. and Bayley, H.S., 1982. Bioenergetics of salmonids fishes: energy intake expenditure and productivity. Comp. Biochem. Physiol. 73: 25-41.
- Chou, B.S. and Shiao, S.Y. 1996. Optimal dietary lipid level for growth of juvenile hybrid tilapia, *Oreochromis niloticus* x *Oreochromis aureus*. Aquaculture 143: 185-195.
- Chung, T.K. 2002. How to get the best out of phytase. Feed Mix. 10: 27-29.

- Clark, J.S. 1989. The mineral requirement of finfish: A review. In: Proceeding of the People Republic of China Aquaculture and Feed Workshop. September 17-30.
- Cole, S. 2002. Comparative feed enzymology –the single currency of enzyme activity. Feed mix. 10: 32-34.
- Coloso, R.M., King, K., Fletcher J.W., Hendrix, M.A., Subramanyam, M., Weis, P. and Ferraris, R.P. 2003. Phosphorus utilization in rainbow trout (*Oncorhynchus mykiss*) fed practical diets and its consequences on effluent phosphorus levels. Aquaculture 220: 801–820.
- Cosgrove, D.J. 1980. Inositol Phosphates. Their chemistry, biochemistry and physiology. Amsterdam: Elsevier Science Publishers.
- Davis, D.A. and Gatlin, D.M. 1991. Dietary mineral requirements of fish and shrimp. In: (eds. Akiyama, D.M. and Tan, R.K.H.), Proceedings of the Aquaculture Feed Processing and Nutrition Workshop. American Soybean Association, Singapore.
- De Silva, S.S. and Anderson, T.A. 1995. Fish Nutrition in Aquaculture. London: Chapman and Hall.
- [De Silva, S.S. and Radampola, K. 1990. Effect of dietary protein level on the reproductive performance of \*Oreochromis niloticus\*. Proceedings of the Second Asian Fisheries Forum, Asian Fisheries Society, Manila, Philippines.](#)
- Debnath, D., Pal, A.K., Sahu, N.P., Jain, K.K., Yengkokpam, S. and Mukherjee, S.C. 2005a. Effect of dietary microbial phytase supplementation on growth and nutrient digestibility of *Pangasius pangasius* (Hamilton) fingerlings. Aquacult. Res. 36: 180-187.
- Debnath, D., Sahu, N.P., Pal, A.K., Jain, K.K., Yengkokpam, S. and Mukherjee, S.C. 2005b. Mineral status of *Pangasius pangasius* (Hamilton) fingerlings in relation to supplemental phytase: absorption, whole-body and bone mineral content. Aquacult. Res. 36: 326-335.

- Degani, G., Viola, S. and Yehuda, Y. 1997. Apparent digestibility of protein and carbohydrate feed ingredients for adult tilapia (*Oreochromis aureus* x *O. niloticus*). *Isr. J. Aquacult./Bamidgeh* 49: 115-123.
- Dey, M.M., Rab, M.A., Paraguas, F.J., Piumsombun, S., Bhatta, R., Alam, F.Md., Koeshendrajana, S. and Ahmed, M. 2002. Socioeconomics of Freshwater Fish Farming in Asian . *in* Semi-annual Progress Report Strategies and Option for Increasing and Sustaining Fisheries and Aquaculture Production to Benefit Poor Households in Asian. March - August 2002. Available for [http://www.worldfishcenter.org/demandsupply/inception\\_reportaug02/ir\\_aug02\\_appendix2.asp#1](http://www.worldfishcenter.org/demandsupply/inception_reportaug02/ir_aug02_appendix2.asp#1). accessed on 1 September 2006.
- Dey, P. M. and Harborne, J. B. 1990. Methods in Plant Biochemistry. Vol 2. Carbohydrates. London: Academic Press.
- Duncan, D.B. 1955. Multiple-rage and multiple F tests. *Biometrics* 11: 1-42.
- Dupree, H.K. and Sneed, K.P. 1966. Response of channel catfish fingerling to different levels of major nutrients in purified diets. U.S. Bureau of Sports Fish and Wildlife Tech. Pap. No. 9.
- Ellestad, L.E., Dahl, G., Angel, R. and Soares, Jr.J.H. 2003. The effect of exogenously administered recombinant bovine somatotropin on intestinal phytase activity and in vivo phytate hydrolysis in hybrid striped bass *Morone chrysops* × *M. saxatilis*. *Aquacul. Nutr.* 9: 327-336.
- El-Saidy, D.M.S. and Gaber, M.M.A. 2003. Replacement of fish meal with a mixture of different plant protein sources in juvenile Nile tilapia, *Oreochromis niloticus* (L.) dites. *Aquaculf. Res.* 34: 1119-1127.
- El-Sayed, A. M., Mansour, C. R. and Ezzat, A.A. 2003. Effects of dietary protein levels on spawning performance of Nile tilapia (*Oreochromis niloticus*) broodstock reared at different water salinities. *Aquaculture* 220: 619-632.
- El-Sayed, A.M. and Teshima, S. 1991. Tilapia nutrition in aquaculture. *Review in Aquatic Sciences*. 5: 247-265.

- El-Sayed, A.M., Ezzat, A. and Mansour, C.R. 2004. Nutritional requirements of red tilapia. *J. Aquacult. Trop.* 19: 179-190.
- Eya, J.C. and Lovell, R.T. 1997a. Net absorption of dietary phosphorus from various inorganic sources and effect of fungal phytase on net absorption of plant phosphorus by channel catfish *Ictalurus punctatus*. *J. World Aquacult. Soc.* 28: 386-391.
- Eya, J.C. and Lovell, R.T. 1997b. Available phosphorus requirement of food-size channel catfish (*Ictalurus punctatus*) fed practical diets in ponds. *Aquaculture* 154: 283-291.
- FAO.1983. Fish Feeds and Feeding in Developing Countries - An Interim Report on the ADCP Feed Development Program. ADCP/REP/83/18. Rome: The United Nations.
- FAO. 1994. Nutrition of fish and crustaceans: a laboratory manual. GCP/RLA/102/ITA. Rome: The United Nations.
- Fitzsimmons, K. 2005. Tilapia nutrition overview feeding shrimp' s partner in polyculture. *Feed international.* 3: 24-27.
- Fitzsimmons, K. 1997. Introduction to Tilapia Aquaculture Conference Proceedings. In *Tilapia Aquaculture* (K. Fitzsimmons ed.). Proceedings from the Fourth International Symposium on Tilapia in Aquaculture, Vol. 1. Northeast Regional Agricultural Engineering Service: Cooperative Extension, Ithaca, New York.
- Fontainhas-Fernandes, A., Gomes, E., Reis-Henriques, M.A. and Coimbra, J. 1999. Replacement of fish meal by plant proteins in the diet of Nile tilapia: digestibility and growth performance. *Aquacult. Int.* 7: 57-67.
- Forster, I. 1999. A note on the method of calculating digestibility coefficients of nutrients provided by single ingredients to feeds of aquatic animals. *Aquacul. Nutr.* 5: 143-145.

- Francis, G., Makkar, H.P.S. and Becker, K. 2001. Antinutritional factors present in plant-derived alternate fish feed ingredients and their effect in fish. *Aquaculture* 199: 197-227.
- Furukawa, A. and Tsukahara, H. 1966. On the acid digestion for the determination of chromic oxide as an index substance in the study of digestibility of fish feed. *Bull. Jpn. Soc. Sci. Fish.* 32: 502-506.
- Furuya, W. M., Gonçalves, G. S., Furuya, V. R. B. and Hayashi, C. 2001. Phytase as feeding for Nile tilapia (*Oreochromis niloticus*) performance and digestibility. *Rev. Bras. Zootec.* 30: 924-929.
- Getachew, T. 1989. Stomach pH, feeding rhythm and ingestion rate in *Oreochromis niloticus* L. (Pisces: Cichlidae) in Lake Awasa, Ethiopia. *J. Hydrobiologia* 174: 43-48.
- Gifford, S.R. and Clydesdale, F.M. 1990. Interactions among calcium, zinc and phytase with protein sources. *J. Food Sci.* 55: 1720-1724.
- Goddard, J.S. and Mclean, E. 2001. Acid-insoluble ash as an inert reference material for digestibility studies in tilapia, *Oreochromis aureus*. *Aquaculture* 194: 93-98.
- Grisdale, H.B. and Helland, S.J. 1998. Macronutrient utilization by Atlantic halibut (*Hippoglossus hippoglossus*): diet digestibility and growth of 1 kg fish. *Aquaculture* 166: 57-65.
- Hafedh, Y.S.A. 1999. Effects of dietary protein on growth and body composition of Nile tilapia, *Oreochromis niloticus* L. *Aquaculf. Res.* 30: 385-393.
- Hajen, W.E. Beames, R.M., Higgs, D.A. and Dosanjh, B.S. 1993. Digestibility of various feedstuffs by post-juvenile chinook salmon (*Oncorhynchus tshawytscha*) in sea water. 1. Validation of technique. *Aquaculture* 112: 321-332.
- Hanley, F. 1987. The digestibility of foodstuffs and the effects of feeding selectivity on digestibility determinations in tilapia, *Oreochromis niloticus* (L). *Aquaculture* 66: 163-179.

- Hardy, R.W. 2000. New developments in aquatic feed ingredients, and potential of enzyme supplements. In: Cruz -Suárez, L.E., Ricque-Marie, D., Tapia-Salazar, M., Olvera-Novoa, M.A. y Civera-Cerecedo, R., (eds.). Avances en Nutrición Acuícola V. Memorias del V Simposium Internacional de Nutrición Acuícola. 19-22 Noviembre, 2000. Mérida, Yucatán, Mexico.
- Hardy, R.W. and Gatlin, D. 2002. Nutritional strategies to reduce nutrient losses in intensive aquaculture. In Avances en Nutriction Acuicola VI Simposium Internacional de Nutricion Acuicola. Cancum. Quinta Roo. Mexico, 3 al 6 de Septiembre del 2002.
- Haylor, G.S., Beveridge, M.C.M. and Jauncey, K. 1988. Phosphorus nutrition of juvenile *Oreochromis niloticus*. In The Second International Symposium on Tilapia in Aquaculture. ICLARM Conference Proceedings 15, 623 pp. Department of Fisheries, Bangkok and ICLARM, Manila.
- Hua, K. and Bureau, D.P. 2006. Modelling digestible phosphorus content of salmonid fish feeds. Aquaculture 254: 455-465.
- Jackson, L.S., Li, M.H. and Robinson, E.H. 1996. Use of microbial phytase in channel catfish. *Ictalurus punctatus*. J. World Aquacult. Soc. 27: 309-313.
- Jahan, P., Watanabe, T., Satoh, S. and Kiron, V. 2003. Reduction in elemental waste loading from Commercial Carp feeds by manipulating the dietary phosphorus levels. Fish. Sci. 69: 58-65.
- Jobling, M. 1994. Fish Bioenergetics. New York: Chapman and Hall.
- Jongbloed, A.W., Kemme, P.A. and Mroz, Z. 1993. The role of microbial phytase in pig production. In: Enzymes in Animal Nutrition.( Wenk, C. and Booessinger, M. eds). Proceedings of the 1<sup>st</sup> Symposium. Switzerland.
- Ketola, H.G. 1975. Requirement of Atlantic salmon for dietary phosphorus. Trans. Am. Fish. Soc. 104: 548-551.
- Kim, J.D., Kim, K.S., Song, J.S., Lee, J.Y. and Jeong, K.S. 1998. Optimum level of dietary monocalcium phosphate based on growth and phosphorus excretion of mirror carp, *Cyprinus carpid*. Aquaculture 161: 337-344.

- Konietzny, U. and Greiner, R. 2002. Molecular and catalytic properties of phytase-degrading enzyme (phytases). *Int. J. Food Sci. Technol.* 37: 791-812.
- Köprüçü, K. and Özdemir, Y. 2005. Apparent digestibility of selected feed ingredients for Nile tilapia (*Oreochromis niloticus*). *Aquaculture* 250: 308-316.
- Lall, S.P. 1991. Digestibility, metabolism and excretion of dietary phosphorus in fish. In Nutritional Strategies and Aquaculture Waste. Proceedings of the First International Symposium on Nutritional Strategies in Management of Aquaculture Waste. University of Guelph, Guelph Ontario, Canada, pp. 21-36. 1990.
- Lall, S.P. 2002. The minerals. In: Fish Nutrition, 3<sup>rd</sup> (Halver, J.E. and Hardy, R.W. eds.). San Diego: CA Academic Press.
- Lanari, D., Agaro, E.D. and Turri, C. 1998. Use of nonlinear regression to evaluate the effects of phytase enzyme treatment of plant protein diets for rainbow trout *Oncorhynchus mykiss*. *Aquaculture* 161: 345-356.
- Larsen, H.N. and Snieszko, S.F. 1961. Modification of microhematocrit technique with trout blood. *Trans. Amer. Fish. Soc.* 90: 139-142.
- Laws, E.A. 2000. Aquatic Pollution An Introduction Text 3<sup>th</sup> : New York: John Wiley & Sons, Inc.
- Liebert, F. and Portz, L. 2005. Nutrient utilization of Nile tilapia *Oreochromis niloticus* fed plant based low phosphorus diets supplemented with graded levels of different sources of microbial phytase. *Aquaculture* 248: 111-119.
- Liener, I.E. 1994. Implications of antinutritional components in soybean foods. *Crit. Rev. Food Sci. Nutr.* 34: 31-67.
- Lorico-Qucrijcro, B. and Chiu, Y.N. 1989. Protein digestibility studies on *Oreochromis niloticus* using chromic oxide indicator. *Asian Fish. Sci.* 2: 177-191.
- Lovell, R.T. 2002. Various types of meters and transmitters for use in aquaculture, their proper placement, maintenance and operation Proceedings of the 2<sup>nd</sup> International Conference on Recirculation Aquaculture.

- Lovell, R.T. 1978. Dietary phosphorus requirement of channel catfish. Trans. Am. Fish. Soc. 107: 617-621.
- Lovell, R.T. 1989. Nutrition and feeding of fish. New York: Van Nostrand Reinhold.
- Lovell, R.T. 1993. Nutritional value of fish: Scientific status summary. Food Tech. 50: 20-30.
- [Lovell, R.T. 1998. Nutrition and Feeding of Fish 2<sup>nd</sup>. Massachusetts: Kluwer Academic publishers.](#)
- Lovell, R.T. and Limsuwan, T. 1982. Intestinal synthesis and dietary non essentiality of vitamin B<sub>12</sub> for Tilapia nilotica. Trans. Amer. Fish. Soc. 111: 485-490.
- Luquet, P. 1989. Practical consideration on the protein nutrition and feeding of tilapia. Aquat. Living. Resour. 2: 99-104.
- Maina, J.G., Beames, R.M., Higgs, D., Mbugua, P.N., Iwama, G. and Kisia, S.M. 2002. Digestibility and feeding value of some feed ingredients fed to tilapia *Oreochromis niloticus*(L.). Aquaculf. Res. 33: 853-862.
- Miranda, E.C., Pezzato, A.C., Pezzato, L.E. and Furuya, W.M. 2000. Apparent phosphorus availability in food for the Nile Tilapia (*Oreochromis niloticus*). Acta. Sci. 22: 669-675.
- Morales, A.E., Cardenete, G., Sanz, A. and de la Higuera, M. 1999. Re-evaluation of crude fibre and acid-insoluble ash as inert markers, alternative to chromic oxide, in digestibility studies with rainbow trout (*Oncorhynchus mykiss*). Aquaculture 179: 71-79.
- Moreau, Y. 1996. Measurement of the apparent digestibility coefficients for *Oreochromis niloticus* of agro-industrial by-products available in Cote d'Ivoire. In: Pullin, R.S.V., Lazard, J., Legendre, M., Amon Kottias, J.B., Pauly, D. (eds.). The Third International Symposium on Tilapia in Aquaculture. ICLARM Conference Proceedings No. 41, ICLARM, Manila, Philippines.

- Mwachireya, S.A., Beames, R.M., Higgs, D.A., and Dosanjh, B.S. 1999. Digestibility of canola protein products derived from the physical, enzymatic and chemical processing of commercial canola meal in rainbow trout *Oncorhynchus mykiss* (Walbaum) held in fresh water. Aquacul. Nutr. 5: 73–82.
- Nakamura, Y. 1985. Sodium-dependent absorption of inorganic phosphate by the carp intestine. Comp. Biochem. Physiol. 80A: 437-439.
- NRC. 1993. Nutrient Requirement of Fish. Washington, D.C.: National Academy Press.
- Nys, Y., Frapin, D. and Pointillart, A., 1996. Occurrence of phytase in plants, animals and microorganisms. In: Phytase in Animal Nutrition and Waste Management: A BASF Reference Manual 1996. New Jersey: BASF Corporation.
- Ogino, C., and Takeda, H. 1978. Requirement of rainbow trout for dietary calcium and phosphorus in carp and rainbow trout. Bull. Jpn. Soc. Sci. Fish. 45: 1527-1532.
- Ogino, C., Takeuchi, L., Takeda, H. and Watanabe, T. 1979. Availability of dietary phosphorus in carp and rainbow trout. Bull. Jpn. Soc. Sci. Fish. 45: 1538-1553.
- Ogunji, J.O. and Wirth, M. 2002. Influence of dietary protein deficiency on amino acid and fatty acid composition in tilapia, *Oreochromis niloticus*, fingerlings .Isr. J. Aquacult. /Bamidgeh. 54: 64-72.
- Ogunji, J.O., Wirth, M. and Rennert, B. 2002. Assessing the dietary amino acid requirement of tilapia *Oreochromis niloticus* fingerlings. In Challenges to organic farming and sustainable land use in the tropics and subtropics, Deutscher Tropentag, Witzenhausen, 9-11 October 2002.
- Phromkunthong, W. and Gabaudan, J. 2006. Use of microbial phytase to replace inorganic phosphorus. Songklanakarin J. Sci. Tech. 28 (4): 731-743.
- Phromkunthong, W. and Udom, U. 2006. Phosphorus requirement in sex-reversed red tilapia. In XII International Symposium on Fish Nutrition & Feeding, Biarritz, France, May 28<sup>th</sup> - June 1<sup>st</sup> 2006.

- Phromkunthong, W., Gabaudan, J and Musakopat, A. 2005. Use of microbial phytase replace inorganic phosphorus in sex-reversed red tilapia *Oreochromis niloticus* (Linnaeus) x *Oreochromis mosambicus* (Peters). Department of Aquatic Science, Faculty of Natural Resources, Prince of Songkla University, Songkhla, Thailand.
- Phromkunthong, W., Yangthong, M., Supamataya, K. and Nakachart, D. 2004. Effects of phytase enzyme and inorganic phosphate on the utilization of phosphorus in sex-reversed red tilapia *Oreochromis niloticus* (Linn.). Songklanakarin J. Sci. Tech. 26 (2): 181-195.
- Piumsombun, S. 2003. The impact of international fish trade on food security in Thailand. In Report of The Expert Consultation on International Fish Trade and Food Security. 27-30 January 2003. Casablanca. Morocco: FAO Fisheries Report.
- Popma, T.J. 1982. Digestibility of selected feedstuffs and naturally occurring algae by tilapia. PhD Dissertation, Auburn University.
- Portz , L. and Liebert, F. 2004. Growth, nutrient utilization and parameters of mineral metabolism in Nile tilapia *Oreochromis niloticus* (Linnaeus, 1758) fed plant-based diets with graded levels of microbial phytase. J. Anim. Physiol. a. Anim. Nutr. 88: 311–320.
- Powers Hughes, K. and Soares Jr.J.H., 1998. Efficacy of phytase on phosphorus utilization in practical diets fed to striped bass *Morone saxatilis*. Aquacult. Nutr. 4: 133–140.
- Rich, M., Trottier, N.L., KU, P.K. and Garling, D.L. 2001. Apparent digestibility of crude protein and apparent availability of individual amino acids in tilapia (*Oreochromis niloticus*) fed phytase pretreated soybean meal diets. Fish- Physiology-and-Biochemistry 25: 181-194.
- Riche, M. and Brown, P.B. 1996. Availability of phosphorus from feedstuffs fed to rainbow trout (*Oncorhynchus mykiss*). Aquaculture 142: 269-282.

- Robinson, E.H. and Wilson, R.P. 1985. Nutrition and feeding. In: Tucker, C.S. (ed.), Channel Catfish Culture. Development in Aquaculture and Fisheries Science 15: 323 – 404.
- Robinson, E.H., Labomascus, D., Brown, P.B. and Lanton, T.L. 1987. Dietary Calcium and Phosphorus Requirement of *Oreochromis aureus* Reared in Calcium-Free Water. Aquaculture 64: 267-276.
- Rodehutscord, M. and Pfeffer, E. 1995. Effects of supplemental microbial phytase on phosphorus digestibility and utilization in rainbow trout (*Oncorhynchus mykiss*). Wat. Sci. Techn. 31: 143-147.
- Rodehutscord, M., Gregus, Z. and Pfeffer, E. 2000. Effect of phosphorus intake on faecal and non faecal phosphorus excretion in rainbow trout (*Oncorhynchus mykiss*) and the consequences for comparative phosphorus availability studies. Aquaculture 188: 383–398.
- Roy, P.K. and Lall, S.P. 2003. Dietary phosphorus requirement of juvenile haddock (*Melanogrammus aeglefinus* L.). Aquaculture 221: 451–468.
- Saijjadi, M. and Carter, C.G. 2004. Dietay phytase supplementation and the utilization of phosphorus by Atlantic salmon (*Salmo salar* L.) fed a canola-meal-based diet. Aquaculture 240: 417-431.
- Sales, J. and Janssens, GPJ. 2003. Nutrient requirements of ornamental fish .Aquat. Living Resour. 16: 533-540.
- Santiago, C.B. and Lovell, R.T. 1988. Amino acid requirement for growth of Nile tilapia. J. Nutr. 118: 1540-1546.
- Satoh, S., Porn-Ngam, N.K., Takeuchi, T. and Watanabe, T. 1996. Influence of dietary phosphorus levels on growth and mineral availability in rainbow trout. Fish. Sci. 62: 483–487.
- Schafer, A., Koppe, W.M., Meyer-Burgdorff, K.H. and Gunther, K.D. 1995. Effects of a microbial phytase on the utilization of native phosphorus by carp in a diet based on soybean meal. Wat. Sci. Techn. 31: 149-155.

- Shiau, S.Y. 2002 . Tilapia *Oreochromis* spp. In Nutrient Requirements and Feeding of Finfish for Aquaculture. (Webster, C.D. and Lim, C. eds), New York: CAB International Publishing.
- Shiau, S.Y. and Peng, C.Y. 1993. Protein sparing effect by carbohydrates in diets for tilapia, *Oreochromis niloticus* x *O. aureus*. Aquaculture 117: 327-334.
- Shiau, S.Y., and Liang, H.S. 1995. Carbohydrate utilization and digestibility by tilapia, *Oreochromis niloticus* x *O. aureus*, are affected by chromic oxide inclusion in the diet. J. Nutr. 125: 976-982.
- Storebakken, T., Kvien, I. S., Shearer, K. D., Grisdale-Helland, B. G. and Helland, S. J. 1999. Estimation of intestinal evacuation rate in Atlantic salmon (*Salmo salar*) using inert markers and collection of feces by sieving: evacuation of diets with fish meal, soybean meal or bacterial protein. Aquaculture 172: 291-299.
- Storebakken, T., Shearer, K.D. and Roem, A.J. 1998. Availability of protein, phosphorus and other elements in fish meal, soy-protein concentrate and phytase-treated soy-protein-concentrate-based diets to Atlantic salmon, *Salmo salar*. Aquaculture 161: 365-379.
- Sugiura, S.H., Babbitt, J.K., Dong, F.M. and Hardy, R.W. 2000. Utilization of fish and animal by product meals in low-pollution feeds for rainbow trout *Oncorhynchus mykiss* (Walbaum). Aquaculf. Res. 31: 585-593.
- Sugiura, S.H., Gabaudan, J., Dong, F.M., and Hardy, R.W. 2001. Dietary microbial phytase supplementation and the utilization of phosphorus, trace minerals and protein by rainbow trout [*Oncorhynchus mykiss* (Walbaum) ] fed soybean meal-based diets. Aquaculf. Res. 32: 583-592.
- Sugiura, S.H., Raboy, V., Young, K.A., Dong, F.M. and Hardy, R.W. 1999. Availability of phosphorus and trace elements in low-phytate varieties of barley and corn for rainbow trout (*Oncorhynchus mykiss*). Aquaculture 170: 285–296.
- Uhlig, H. 1998. Industrial enzyme and their application. New York: John Wiley & Sons, Inc.

- Tacon, A.G.J. and Forster, I.P. 2003. Aquafeeds and the environment: policy implications. *Aquaculture* 226: 181-189.
- Tacon, A.G.J. and Rodrigues, A.M.P. 1984. Comparison of chromic oxide, crude fibre, polyethylene and acid-insoluble ash as dietary markers for the estimation of apparent digestibility coefficients in rainbow trout. *Aquaculture* 43: 391-399.
- Thiessen, D. Campbell, G. and Adelizi, P. 2003. Digestibility and growth performance of juvenile rainbow trout (*Oncorhynchus mykiss*) fed with pea and canola products. *Aquacul. Nutr.* 9: 67-75.
- Twibell, R.G., and Brown, P.B. 1998. Optimal dietary protein concentration for hybrid tilapia *Oreochromis niloticus* x *O. aureus* fed all-plant diets. *J. World Aquacult. Soc.* 29: 9-16.
- Vielma, J. and Lall, S.P. 1998. Phosphorus utilization by Atlantic salmon (*Salmo salar*) reared in freshwater is not influenced by higher dietary calcium intake. *Aquaculture* 160: 117-128.
- Vielma, J., Lall, S.P., Koskela, J., Schoner, F.-J. and Mattila, P. 1998. Effects of dietary phytase and cholecalciferol on phosphorus bioavailability in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 163: 309-323.
- Vielma, J., Makinen, T., Ekholm, P. and Koskela, J. 2000. Influence of dietary soy and phytase levels on performance and body composition of large rainbow trout (*Oncorhynchus mykiss*) and algal availability of phosphorus load. *Aquaculture* 183: 349-362.
- Vielma, J. and Ruohone, K. 2002. Dephytinization of two soy proteins increases phosphorus and protein utilization by rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 204: 145-156.
- Vielma, J., Ruohonen, K., Gabaudan, J. and Voge, K. 2004. Top-spraying soybean meal-based diets with phytase improves protein and mineral digestibilities but not lysine utilization in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquaculf. Res.* 35: 955-964.

- Vielma, J., Ruohonen, K. and Peisker, M. 2002. Dephytinization of two soy proteins increases phosphorus and protein utilization by rainbow trout, *Oncorhynchus mykiss*. Aquaculture 204: 145-156.
- Viola, S. and Arieli, Y. 1983. Evaluation of different grains as basic ingredients in complete feeds for carp and tilapia in intensive culture. Isr. J. Aquacult./Bamidgeh. 35: 38-43.
- Vohara, A. and Satyanarayana, T. 2003. Phytase: microbial sources production, purification and potential biotechnological applications. Critical Reviews in Biotechnology. 23: 29-60.
- Wang, H.L., Swain, E.W. and Hesseltine, C.W. 1980. Phytase of molds used in oriental food fermentation. J. Food Sci. 45: 1262-1266.
- Watanabe, T., Satoh, S. and Takeuchi, T. 1988. Availability of minerals in fish meal to fish Asian Fish. Soc. 1: 175-195.
- Watanabe, T., Takeuchi, T., Murakami, A. and Ogino, C. 1980a. The availability to *Tilapia niloticus* of phosphorus in white fishmeal. Bull. Jap. Soc. Sci. Fish. 46: 897-899.
- Watanabe, T., Takeuchi, T., Murakami, A. Nose, T. and Ogino, C. 1980b. Requirement of chum salmon held in freshwater for dietary phosphorus. Bull. Jap. Soc. Sci. Fish. 46: 361-367.
- Watanabe, T., Takeuchi, T., Satoh, S. and Kiron, V. 1996. Digestible crude protein contents in various feedstuffs determined with four freshwater fish species. Fish. Sci. 62: 278-282.
- Wilson, R.P., Robinson, E.H. Gatlin III, D.M. and Poe, W.E. 1982. Dietary phosphorus requirement of channel catfish. J. Nutr. 112: 1197-1202.
- Yone, Y. and Fujii, M. 1975. Studies on Nutrition of red sea bream-XI Effect of  $\Omega 3$  fatty acid supplement in a corn oil diet on growth rate and feed efficiency. Bull. Jap. Soc.sci. Fish 41: 73-77.

- Yoo, G.-Y., Wang, X., Choi, S., Han, K., Kang, J.-C. and Bai, S.C. 2005. Dietary microbial phytase increased the phosphorus digestibility in juvenile Korean rockfish *Sebastes schlegeli* fed diets containing soybean meal. Aquaculture 243: 315-322.
- Zeitoun, I.H., Tack, P.I., Halver, J.E. and Ullrey, D.E. 1973. Influence of salinity on protein requirements of rainbow trout *Salmo gairdneri* fingerling. J. Fish. Res. Board Can. 30: 1867-1873.