

เอกสารอ้างอิง

เกรียงศักดิ์ สายธนู, เกรียงศักดิ์ พูนสุข และ สงคราม เหลืองทองคำ. 2524. การแพร่กระจายของไวรัสโ พาราฮีโมลัสติคัส ในน่านน้ำไทย ผลการสำรวจปี 2521-2524. การสัมมนาครั้งที่ 2 การวิจัยคุณภาพน้ำและทรัพยากรมีชีวิตในน่านน้ำไทย สำนักงานคณะกรรมการวิจัยแห่งชาติ กรุงเทพฯ. หน้า 255-261.

บุญเยี่ยม เกียรติวุฒิ, อรุณ เกียรติวุฒิ และ ศุภกิจ อังคศุภากร 2527. โรคติดต่อระหว่างคนและสัตว์ หน้า 92-96. กรุงเทพฯ:บัณฑิตการพิมพ์.

พรนภัส ประเสริฐไทยเจริญ, ชันยาพร เพิ่มวงศ์ การเฝ้าระวังทางห้องปฏิบัติการของเชื้อแบคทีเรียที่ทำให้เกิดโรคในลำไส้คน จังหวัดอุบลราชธานี เมษายน 2543-มีนาคม 2544. <http://www.moph.go.th/ops/epi/known/lab.html>.

อรษา สุตเธียรกุล. 2541. โรคติดเชื้อ. พิมพ์ครั้งที่ 2. กรุงเทพฯ:บริษัท โฮลิสติก พับลิชชิง จำกัด.

Alouf, J.E. and Freer, J.H. 1999. The Comprehensive Sourcebook of Bacterial Protein Toxins. 2nd ed London:Academic Press.

Altekruse, S.F., Bishop, R.D., Baldy, L.M., Thompson, S.G., Wilson, S.A., Ray, B.J. and Griffin, P.M. 2000. *Vibrio* gastroenteritis in the US Gulf of Mexico region: the role of raw oysters. *Epidemiol. Infect.* 124:489-495.

Baffone, W., Pianehi, A., Bruscolini, F., Barbieri, E. and Citterio, B. 2000. Occurrence and expression of virulence-related properties of *Vibrio* species isolated from widely consumed seafood products. *Int. J. Food Microbiol.* 54:9-18.

- Baba, K., Shirai, H., Terai, A., Kumagai, K., Takeda, Y. and Nishibuchi, M. 1991. Similarity of the *tdh* gene-bearing plasmid of *Vibrio cholerae* non-O1 and *Vibrio parahaemolyticus*. *Microb. Pathog.* 10:61-70.
- Bag, P.K., Nandi, S., Bhadra, R.K., Ramamurthy, T., Bhattacharya, S.K., Nishibuchi, M., Hamabata, T., Yamasaki, S., Takeda, T. and Nair, B.G. 1999. Clonal diversity among recently emerged strains of *Vibrio parahaemolyticus* O3:K6 associated with pandemic spread. *J. Clin. Microbiol.* 37:2354-2357.
- Balows, A., Truper, H.G., Dworkin, M., Harder W. and Schleifer K.H. 1992. *The Prokaryotes* 2nd ed. New York: Springer-Verlag New York Inc.
- Barker, Jr. W.H., and Gangarosa E.J. 1974. Food poisoning due to *Vibrio parahaemolyticus*. *Annu. Rev. Med.* 25:75-81.
- Barnes D.R. 1974. *Invertebrate Zoology*. 3rd ed. Philadelphia: WB. Saunders Company.
- Centers for Disease Control and Prevention. 1999. Outbreak of *Vibrio parahaemolyticus* infection associated with eating raw oysters and clams harvested from Long Island Sound-Connecticut, New Jersey, and New York 1998. *JAMA* 281:603-604.
- Chan, K.Y., Woo, M.L., Lam, L.Y. and French, G.L. 1989. *Vibrio parahaemolyticus* and other halophilic vibrios associated with sea food in Hong Kong. *J. Appl. Bacteriol.* 66:57-64.
- Chen, A., Edgar, D.B. and Trela, J.M. 1976. Deoxyribonucleic acid polymerase from the extreme thermophile *Thermus aquaticus*. *J. Bacteriol.* 127:1550-1557.

- Chiou, C., Hsu, S., Chiu, S., Wang, T. and Chao, C. 2000. *Vibrio parahaemolyticus* serovar O3:K6 as cause of unusually high incidence of food-borne disease outbreaks in Taiwan from 1996 to 1999. *J. Clin. Microbiol.* 38:3981-3986.
- Chowdhury, N.R., Chakraborty, S., Eampokalap, B., Chaicumpa, W., Chongsanguan, M., Moolasart, P., Mitra, R., Ramamurthy, T., Bhattacharya, S.K., Nishibuchi, M., Takeda, Y. and Nair, G.B. 2000a. Clonal dissemination of *Vibrio parahaemolyticus* displaying similar DNA fingerprint but belonging to two different serovars (O3:K6 and O4:K68) in Thailand and India. *Epidemiol. Infect.* 125:17-25.
- Chowdhury, N.R., Chakraborty, S., Ramamurthy, T., Nishibuchi, M., Yamasaki, S., Takeda, Y. and Nair, G.B. 2000b. Molecular evidence of clonal *Vibrio parahaemolyticus* pandemic strains. *Emerg. Infect. Dis.* 6:631-633.
- Cook, D.W., O'Leary, P., Hunsucker, J.C., Sloan, E.M., Bowers, J.C., Blodgett, R.J., and DePaola A. 2002. *Vibrio vulnificus* and *Vibrio parahaemolyticus* in U.S. retail shell oysters : a national survey from June 1998 to July 1999. *J. Food Prot.* 65:79-87.
- Daniels, N.A., MacKinnon, L., Bishop, R., Ahekruse, S., Ray, B., Hammond, R.M., Thompson, S., Wilson, S., Bean, N.H. , Griffin, P. M. and Slutsker, L. 2000. *Vibrio parahaemolyticus* infection in the United States 1973-1998. *J. Infect. Dis.* 181:1661-1666.
- DePaola, A., Hopkins, L.H., Peeler, J.T., Wentz, B. and Mc Phearson, R.M. 1990. Incidence of *Vibrio parahaemolyticus* in United State coastal water and oysters. *Appl. Environ. Microbiol.* 56:2299-2302.

- DiRita , V.J. 1992. Co-ordinate expression of virulence genes by ToxR in *Vibrio cholerae*.
Molecular Microbiol. 6:451-458.
- Echeverria, P., Pitarangsi, C., Eampokalap, B., Vibulbandhitkit, S., Boonthai, P. and Rowe,
B. 1983. A logitudinal study of the prevalence of bacterial enteric pathogens among
adults with diarrhea in Bangkok, Thailand. Diagnostic Microbiol. Infect. Dis. 1:193-
204.
- Fabbri, A., Falsano, L., Frank, C., Donelli, G., Matarrese, P., Raimondi, F., Fasano, A., and
Fiorentini, C. 1999. *Vibrio parahaemolyticus* thermostable direct hemolysin
modulated cytoskeletal organization and homeostasis in intestinal culture cells.
Infect. Immun. 67:1139-1148.
- Farmer, J.J., Hickman-Brenner, F.W. and Kelly, M.T. 1985. Vibrio. In : Lennette, E.H.,
Balows, A., JR., Hausler W.J. and Shadomy, H.J. (eds.). Manual of Clinical
Microbiology p. 282-301. 5th ed. Washington, D.C.: American Society for
Microbiology.
- Ghosh, A.R. and Sehgal, S.C. 1998. Detection of *tdh* and *trh* genes in urea-hydrolysing
environmental isolate of *Vibrio parahaemolyticus* from the Andamans. J. Diarrhoeal
Dis. Res. 16:000-000
- Hackney, C.R., Ray, B. and Speck, M. L. 1980. Incidence of *Vibrio parahaemolyticus* in
and the microbiological quality of seafood in North Carolina. J. Food Prot. 43:769-
773

- Hervio-Health, D., Colwell, R.R., Derrien, A., Robert-Pillot, A., Fournier, J.M. and Pommepuy, M. 2002. Occurrence of pathogenic Vibrios in coastal area of France. *J. Appl. Microbiol.* 92:1123-1135.
- Hickman, F.W., Farmer, J.J., Hollis, D.G., Fanning, G.R., Steigerwalt, A.G., Weaver, R.E. and Brenner, D.J. 1982. Identification of *Vibrio hollisae* sp. nov. from patients with diarrhea. *J. Clin. Microbiol.* 15:395-401.
- Hlady, W.G. 1997. Vibrio infections associated with raw oyster consumption in Florida, 1981-1994. *J. Food Prot.* 60:353-357.
- Honda, T., Goshima, K., Takeda, Y., Sugino, Y. and Miwatani, T. 1976a. Demonstration of the cardiotoxicity of the thermostable direct hemolysin (lethal toxin) produced by *Vibrio parahaemolyticus*. *Infect. Immun.* 13:163-171.
- Honda, S., Goto, I., Minematsu, I., Ikeda, N., Asano, N., Ishibashi, M., Kinoshita, Y., Nishibuchi, M., Honda, T. and Miwatani, T. 1987. Gastroenteritis due to Kanagawa-negative *Vibrio parahaemolyticus*. *Lancet* I:331-332
- Honda, T. and Iida, T. 1993. The pathogenicity of *Vibrio parahaemolyticus* and the role of the thermostable direct hemolysin and related hemolysins *Rev. Med. Microbiol.* 4:106-113
- Honda, T., Ni, Y., Hata, A., Yoh, M., Miwatani, T., Okamoto, T., Goshima, K., Takakura, H., Tsunasawa, S. and Sakiyama, F. 1990. Properties of a hemolysin related to the thermostable direct hemolysin produced by a kanagawa phenomenon negative, clinical isolate of *Vibrio parahaemolyticus*. *Can. J. Microbiol.* 36:395-399.

- Honda, T., Ni, Y. and Miwatani, T. 1988. Purification and characterization of a hemolysin produced by a clinical isolate of Kanagawa phenomenon-negative *Vibrio parahaemolyticus* and related to the thermostable direct hemolysin. *Infect. Immun.* 56:961-965.
- Honda, T., Taga, S., Takeda, T., Hasibuan, M.A., Takeda, Y., and Miwatani, T. 1976b. Identification of lethal toxin with thermostable direct hemolysin produce by *Vibrio parahaemolyticus* and some physico-chemical properties of the purified toxin. *Infect. Immun.* 13:133-139.
- Iguchi, T., Kondo, S. and Hisatsune K. 1995. *Vibrio parahaemolyticus* O serotypes from O1 to O13 all produce R-type lipopolysaccharide: SDS-PAGE and compositional sugar analysis. *FEMS Microbiol. Lett.* 130:287-292.
- Iida, T., Park, K.S. Suthienkul, O., Kozawa, J., Yamaichi, Y., Yamamoto, K. and Honda, T. 1998. Close proximity of the *tdh trh* and *ure* genes on the chromosome of *Vibrio parahaemolyticus*. *Microbiology.* 144:2517-2523.
- Iida, T., Suthienkul, O., Park, K., Tang, G.Q., Yamamoto, R.K., Ishibashi, M., Yamamoto, K. and Honda, T. 1997. Evidence for genetic linkage between the *ure* and *trh* genes in *Vibrio parahaemolyticus* . *J. Med. Microbiol.* 46:639-645.
- Iida, T. and Yamamoto, K. 1990 Cloning and expression of two genes encoding highly homologous hemolysins from a Kanagawa phenomenon-positive *Vibrio parahaemolyticus* T4750 strain.

- Johnton, M.D. and Brown, M.H. 2002. An investigation into the changed physiological state of *Vibrio* bacteria as a survival mechanism in response to cold temperatures and studies on their sensitivity to heating and freezing. *J. Appl. Microbiol.* 92:1066-1077.
- Kaneko, T. and Cowell, R.R. 1973. Ecology of *Vibrio parahaemolyticus* in Chesapeake Bay. *J. Bacteriol.* 113:24-32.
- Kaysner, C.A., Abeyta, C. J.R., Stott, R.F., Lilja, J.L. and Wekell, M.M. 1990. Incidence of urea-hydrolyzing *Vibrio parahaemolyticus* in Willapa Bay, Washington. *Appl. Environ. Microbiol.* 56:904-907.
- Kelly, M.T. and Stroh E.M.D. 1989. Urease-positive, Kanagawa-negative *Vibrio parahaemolyticus* from patients and environment in the Pacific North West. *J. Clin. Microbiol.* 27:2820-2822.
- Kim, T.B., Okuda, J., Matsumoto, C., Takahashi, N., Hashimoto, S. and Nishibuchi, M. 1999. Identification of *Vibrio parahaemolyticus* strains at the species level by PCR targeted to the *toxR* gene. *J. Clin. Microbiol.* 37:1173-1177
- Kishishita, M., Matsuoka, N., Kumagai, K., Yamasaki, S., Takeda, Y. and Nishibushi, M. 1992. Sequence variation in the thermostable direct hemolysin-related hemolysin (*trh*) gene of *Vibrio parahaemolyticus*. *Appl. Environ. Microbiol.* 58:2449-2457.
- Lee, C., Chen, L.H., Liu, M.L. and Su, Y.C. 1992. Use of an oligonucleotide probe to detect *Vibrio parahaemolyticus* in artificially contaminate oysters. *Appl. Environ. Microbiol.* 58:3419-3422.

- Lee, J.V. 1990. *Vibrio*, *Aeromonas* and *Plesiomonas* In: Parker, M.T. and Duerden, B.I. (eds.). Principles of Bacteriology, Virology and Immunity, 8th ed. Vol II. p. 514-527. Philadelphia: B.C. Decker.
- Lee, S.E., Shin, S.H., Kim, S.Y., Kim, Y.R., Shin, D. H., Chung, S.S., Lee, Z.H., Lee, J.Y., Jeong, K.C., Choi, S.H. and Rhee, J.H. 2000. *Vibrio vulnificus* has the transmembrane transcription activator ToxRS stimulating the expression of the hemolysin gene *vvha*. J. Bacteriol. 182:3405-3415.
- Lin, Z., Kumagai, K., Baba, K., Mekalanos, J.J. and Nishibuchi, M. 1993. *Vibrio parahaemolyticus* has a homolog of the *Vibrio cholerae* *toxRS* operon that mediates environment induced regulation of the thermostable direct hemolysin gene. J. Bacteriol. 175:3844-3855.
- Lowry, P.W., McFarland, L.M., Pelitier, B.H., Roberts, N. C., Bradford, H.B., Herndon, J.L., Stoup, D.F., Mathison, J.B., Blake, P.A. and Gunn, R.A. 1989. *Vibrio* gastroenteritis in Louisiana: a prospective study among attendees of scientific congress in New Orleans. J. Infect. Dis. 160:978-984.
- Marshall, S., Clark, C.G., Wang, G., Mulvey, M., Kelly, M.T. and Johnson, W. M. 1999. Comparison of molecular methods for typing *Vibrio parahaemolyticus*. J. Clin. Microbiol. 37:2473-2478.
- Matsumoto, C., Okuda, J., Ishibashi, M., Iwanaga, M., Garg P., Ramamurthy, T., Wong, H., Depaola, A., Kim, Y.B., Albert, J.M. and Nishibuchi, M. 2000. Pandemic spread of an O3:K6 clone of *Vibrio parahaemolyticus* and emergence of related strains

- evidenced by arbitrarily primed and *toxRS* sequence analyses. J. Clin. Microbiol. 38:578-585
- Miller, V.L., and Mekalanos J.J. 1988. A novel suicide vector and its use in construction of insertion mutation: osmoregulation of outer membrane proteins and virulence determinants in *Vibrio cholerae* requires *toxR*. J. Bacteriol. 170:2575-2583.
- Miller, V.L., DiRita, V.J. and Mekalanos, J.J. 1989. Identification of *toxS*, a regulatory gene whose product enhances *ToxR*-mediated activation of the cholera promoter. J. Bacteriol. 171:1288-1293.
- Miller, V.L., Taylor, R.K. and Mekalanos, J.J. 1987. Cholerae toxin transcriptional activator *ToxR* is a transmembrane DNA binding protein. Cell. 48:271-279.
- Ming, X., Yamamoto, K. and Honda, T. 1994. Construction and characterization of an isogenic mutant of *Vibrio parahaemolyticus* having a deletion in the thermostable direct hemolysin-related hemolysin gene (*trh*). J. Bacteriol. 176:4757-4760.
- Miwatani, T. and Takeda, Y. 1976. *Vibrio parahaemolyticus* a Causative Bacterium of Food Poisoning. Tokyo:Saikon Publishing Co., LTD.
- Miyamoto, Y., Kato, T., Obara, Y., Akiyama, S., Takizawa, K. and Yamai, S. 1969. In vitro hemolytic characteristic of *Vibrio parahaemolyticus*: its close correlation with human pathogenicity. J. Bacteriol. 100:1147-1149.
- Miyamoto, Y., Obara, Y., Nikkawa, T., Yamai, S., Kato, T., Yamada, Y. and Ohashi, M. 1980. Simplified purification and biophysicochemical characteristics of Kanagawa

- phenomenon-associated hemolysin of *Vibrio parahaemolyticus*. Infect. Immun. 28:567-576.
- Myatt, D.C. and Davis, G.H. 1989. Isolation of medically significant *Vibrio* species from riverine sources in Southeast Queensland. Microbiol. 60:111-123.
- Nagayama, K., Yamamoto, K., Mitawani, T. and Honda, T. 1995. Characterisation of a haemolysin related to Vp-TDH produced by a Kanagawa phenomenon-negative clinical isolate of *Vibrio parahaemolyticus* J. Med. Microbiol. 42:83-90.
- Nishibuchi, M., Doke, S., Toizumi, S., Umeda, T., Yoh, M. and Miwatani, T. 1988. Isolation from a coastal fish of *Vibrio hollisae* capable of producing a hemolysin similar to the thermostable direct hemolysin of *Vibrio parahaemolyticus*. Appl. Environ. Microbiol. 54:2144-2146.
- Nishibuchi, M., Hill, W.E., Zon, G., Payne, W.L. and Kaper J.B. 1986. Synthetic oligodeoxyribonucleotide probes to detect Kanagawa phenomenon-positive *Vibrio parahaemolyticus*. J. Clin. Microbiol. 23:1091-1095.
- Nishibuchi, M., Kumagai, K. and Kaper, J.B. 1991. Contribution of the *tdh1* gene of Kanagawa phenomenon-positive *Vibrio parahaemolyticus* to production of extracellular thermostable direct hemolysin. Microb. Pathog. 11:453-460.
- Nishibuchi, M., Taniguchi, T., Misawa, T., Khaeomane-IAM, V., Honda, T. and Miwatani, T. 1989. Cloning and nucleotide sequence of the genetic encoding the hemolysin related to the thermostable direct hemolysin of *Vibrio parahaemolyticus*. Infect. Immun. 57:2691-2697.

- Nishibuchi, M., and Kaper, J.B. 1990. Duplication and variation of the thermostable direct hemolysin (*tdh*) gene in *Vibrio parahaemolyticus*. *Mol. Microbiol.* 4:87-99.
- Nishibuchi, M., and Kaper, J.B. 1995. Thermostable direct hemolysin gene of *Vibrio parahaemolyticus* : a virulence gene acquired by a marine bacterium. *Infect. Immun.* 64:2093-2099.
- Okitsu, T., Osawa, R., Pornruangwong, S., and Yama, S. 1997. Urea hydrolysis and suppressed production of thermostable direct hemolysin (TDH) by *Vibrio parahaemolyticus* associated with presence of TDH-related hemolysin genes. *Curr. Microbiol.* 34:314-317.
- Okuda, J., Ishibashi, M., Hayakawa, E., Nishino, T., Takeda, Y., Mukhopadhyay, A.K., Garg, S., Bhattacharya, S.K., Nair, G.B. and Nishibuchi, M. 1997a. Emergence of unique O3:K6 clone of *Vibrio parahaemolyticus* in Calcutta, India, and isolation of strains from the same clonal group from Southeast Asian travelers arriving in Japan. *J. Clin. Microbiol.* 35:3150-3155.
- Okuda, J. and Nishibuchi, M. 1998. Manifestation of the Kanagawa phenomenon the virulence-associated phenotype of *Vibrio parahaemolyticus* depend on a particular single base change in the promoter of the thermostable direct hemolysin gene. *Mol. Microbiol.* 30:499-511.
- Okuda, J., Ishibashi, M., Abbott, S.I., Janda, J.M. And Nishibuchi, M. 1997b. Analysis of the thermostable direct hemolysin (*tdh*) and the *tdh*-related hemolysin (*trh*) gene in urease-positive strains of *Vibrio parahaemolyticus* isolated on the west coast of the United States. *J. Clin. Microbiol.* 35:1965-1971.

- Olive, D.M. and Bean, P. 1999. Principle and applications of methods for DNA-based typing of microbial organisms. *J. Clin. Microbiol.* 37:1661-1669
- Oliver, J.D. and Kaper, J.B. 1997. *Vibrio* species. In : Doyle, M.P., Beuchat, L.R. and Montville, T.J. (eds). *Food Microbiology Fundamentals and Frontiers*. P.228-264 Washington, D.C. : American Society for Microbiology.
- Osorio, C.R. and Klose, K.E. 2000. A region of the transmembrane regulatory protein ToxR that tethers the transcriptional activation domain to the cytoplasmic membrane displays wide divergence among *Vibrio* species. *J. Bacteriol.* 182:526-528.
- Raimondi, F., Kao, J.P.Y., Fiorentini, C., Fabbri, A., Donelli, G., Gasparini, N., Rubino, A. and Fasano, A. 2000. Enterotoxicity and cytotoxicity of *Vibrio parahaemolyticus* thermostable direct hemolysin *in vitro* systems. *Infect. Immun.* 68:3180-3185.
- Reich, K. A. and Schoolnik G.K. 1994. The light organ symbiont *Vibrio fischeri* posses a homolog of the *Vibrio cholerae* transmembrane transcriptional activator ToxR. *J. Bacteriol.* 176:3085-3088.
- Sakurai, J., Honda, T., Jinguji, Y., Arita, M. and Miwatani, T. 1976. Cytotoxic effect of the the thermostable direct hemolysin produce by *Vibrio parahaemolyticus* on FL cells. *Infect. Immun.* 13:876-883.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. 1989. *Molecular Cloning: a Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.

- Sanyal, S.C. and Sen, P.C. 1974. Human volunteer study on the pathogenicity of *Vibrio parahaemolyticus*. In : Fujino, T., Sakaguchi, G., Sakazaki, R. and Takeda, Y. (eds.). International Symposium on *Vibrio parahaemolyticus*.p. 227-230. Tokyo:Saikon Publising Co, Ltd.
- Schwartz, D.C. and Cantor, C.R. 1984. Separation of yeast chromosome-sized DNAs by pulse field gel electrophoresis. *Cell*. 37:67-75.
- Sehgal, S.C. and Ghosh, A.R. 1998. Detection of *tdh* and *trh* genes in a urea-hydrolysing environmental isolate of *Vibrio parahaemolyticus* from the Andamans. *J. Diarrhoeal. Dis. Res.* 16:000-000.
- Singh, D.V., Matte, M.H., Matte, G.R., Jiang, S., Sabeen, F., Shukla, B.N., Sanyal, S.C., Huq, A. and Colwell, R.R. 2001. Molecular analysis of *Vibrio cholerae* O1, O139, non-O1 and non-O139 strains: clonal relationships between clinical and environmental isolates. *Appl. Environ. Microbiol.* 67:910:921.
- Suthienkul, O., Ishibashi, M., Iida, T., Nettip, N., Supavej, S., Eampokatap, B., Makino, M. and Honda, T. 1995. Urease production correlates with possession of the *trh* gene in *Vibrio parahaemolyticus* strains isolated in Thailand. *J. Infect. Dis.* 172:1405-1408.
- Tada, J., Ohashi, T., Nishimura, N., Shirasaki, Y., Ozaki, H., Fugushima, S., Takano, J., Nihsibuchi, M. and Takeda, F. 1992. Detection of the thermostable direct hemolysin gene (*tdh*) and the thermostable direct hemolysin-related hemolysin gene (*trh*) of *Vibrio parahaemolyticus* by polymerase chain reaction. *Mol. Cell Probes.* 6:477-487.

- Takahashi, A., Sato, Y., Shiomi, Y., Cantarelli, V.V., Iida, T., Lee, M. and Honda, T., 2000. Mechanism of chloride secretion induced by thermostable direct hemolysin of *Vibrio parahaemolyticus* in human colonic tissue and intestinal epithelial cell line. *J. Med. Microbiol.* 49:801-810.
- TanG, G.Q., Iida, T., Yamamoto, K. and Honda, T. 1995. Ca²⁺ independent cytotoxic of *Vibrio parahaemolyticus* thermostable direct hemolysin (TDH) on intestine 407, a cell derived from human embryonic intestine. *FEMS Microbiol. Lett.* 134:233-238.
- Tangkanakul, W., Tharmaphorhpilas, P., Datapon, D. and Sutantayawalee, S. 2000. Food poisoning outbreak from contaminated fish-balls. *J. Med. Assoc. Thai.* 83:1289-1295.
- Taniguchi, H., Sato, K., Ogawa, M., Udou, T. and Mizuguchi, Y. 1984. Isolation and characterization of filamentous phage ν f 33, specific for *Vibrio parahaemolyticus*. *Microbiol. Immunol.* 28:327-337.
- Terai, A., Shirai, H., Yoshida, O., Takeda, Y. and Nishibuchi, M. 1990. Nucleotide sequence of the thermostable direct hemolysin gene (*tdh* gene) of *Vibrio mimicus* and its evolutionary relationship with the *tdh* gene of *Vibrio parahaemolyticus*. *FEMS Microbiol. Lett.* 71:319-324.
- Terai, A., Baba, K., Shirai, H., Yoshida, O., Takeda, Y. and Nishibuchi, M. 1991. Evidence for insertion sequence-mediated spread of the thermostable direct hemolysin gene among *Vibrio* species. *J. Bacteriol.* 173:5036-5046.

- Thamlikitkul, V. 1990. *Vibrio* bacterium in Siriraj hospital. J. Med. Assoc. Thai. 73:136-139.
- Tomoyasu, T. 1992. Development of the immunomagnetic enrichment method selective for *Vibrio parahaemolyticus* serotype K and its application to food poisoning study. Appl. Environ. Microbiol. 58:4621-4625.
- Vanderzant, C. and Nickelson, R. 1972. Survival of *Vibrio parahaemolyticus* in shrimp tissue under various environmental conditions. Appl. Environ. Microbiol. 23:34-37
- Vuddhakul, V., Chowdhury, A., Loahaprerththisan, V., Pungrasamee, P., Patararungrong, N., Thianmontri, P., Ishibashi, M., Matsumoto, C. and Nishibuchi M. 2000a. Isolation of a pandemic O3:K6 clone of a *Vibrio parahaemolyticus* strain from environmental and clinical sources in Thailand. Appl. Environ. Microbiol. 66:2685-2689.
- Vuddhakul, V., Nakai, T., Matsumoto, C., Oh, T., Nishino, T., Chen, C.H., Nishibuchi, M. and Okuda, J. 2000b. Analysis of *gryB* and *toxR* gene sequences of *Vibrio hollisae* and development of *gryB*- and *toxR*- targeted PCR methods for isolation of *Vibrio hollisae* from the environment and its identification . Appl. Environ. Microbiol. 66:3506-3514.
- Weagant, S.D. and Kaysner, C.A. 1982. The incidence and seasonal distribution of *Yersinia enterocolitica* and *Vibrio parahaemolyticus* in Puget Sound commercial oyster bed. J. Shellfish Res. 2:122-123.

- William, J.G.K., Kubelik, A.R., Livak, K.J., Rafalski, J.A. and Tingey, S.V. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acid Research*. 18:6531-6535.
- Wong, H.C., Chen, M.C., Liu, S.H. and Liu, D.P. 1999. Incidence of highly genetically diversified *Vibrio parahaemolyticus* in seafood imported from Asian countries *Int. Food Microbiol.* 52:181-188.
- Wong, H.C., Chen, L.L. and Yu, C.M. 1994. Survival of psychrotrophic *Vibrio mimicus*, *Vibrio fluvialis* and *Vibrio parahaemolyticus* in culture broth at low temperatures. *J. Food Protect.* 57:607-610.
- Wong, H.C., Lieu, S.H., Wang, T.K., Lee, C.I., Chiou, C.S., Liu, D.P., Nishibuchi, M. and Lee, B.K. 2000. Characteristics of *Vibrio parahaemolyticus* O3:K6 from Asia. *Appl. Environ. Microbiol.* 66:3981-3986.
- Wong, H.C., Lu, K.T., Pan, T.M., Lee, C.I., and Shih D.Y. 1996. Supspecies typing of *Vibrio parahaemolyticus* by Pulse-field gel electrophoresis *J. Clin. Microbiol.* 34:1535-1539.
- Wong, H.C., Ting, S.H. and Shieh, W.R. 1992. Incidence of toxigenic *Vibrios* in foods available in Taiwan. *J. Appl. Bacteriol.* 73:197-202.
- Yamasaki, S., Shirai, H., Takeda, Y. and Nishibuchi, M. 1991. Analysis of the gene of *Vibrio hollisae* encoding the hemolysin similar to the thermostable direct hemolysin of *Vibrio parahaemolyticus*. *FEMS Microbiol. Lett.* 80:259-264.

- Yeung, P.S.M., Hayes, C.M., DePaola, A., Kaysner, A.C., Kornstein, L. and Boor, J. K. 2002. Comparative phenotypic, molecular and virulence characterization of *Vibrio parahaemolyticus* O3:K6 isolates. *Appl. Environ. Microbiol.* 68:2901-2909
- Yoh, M., Honda, T., Miwatani, T. and Nishibuchi, M. 1991. Characterization of thermostable direct hemolysins encoded by four representative *tdh* gene of *Vibrio parahaemolyticus*. *Microb. Pathog.* 10:165-172.