

เอกสารอ้างอิง

- Abraham, W-R., Meyer, H. and Yakimov, M. 1998. Novel glycine containing glucolipids from the alkane using bacterium *Alcanivorax borkumensis*. *Biochim. Biophys. Acta* 1393 : 57-62.
- Batista, S.B., Mounteer, A.H., Amorim, F.R. and Tótoia, M.R. 2006. Isolation and characterization of biosurfactant/bioemulsifier-producing bacteria from petroleum contaminated sites. *Bioresource Technol.*, 97 : 868-875.
- Beal, R. and Betts, W.B. 2000. Role of rhamnolipid biosurfactants in the uptake and mineralization of hexadecane in *Pseudomonas aeruginosa*. *J. Appl. Microbiol.*, 89 : 158-168.
- Bredholt, H., Josefsen, K., Vatland, A., Bruheim, P. and Eimhjellen, K. 1998. Emulsification of crude oil by an alkane-oxidizing *Rhodococcus* species isolated from seawater. *Can. J. Microbiol.*, 44 : 330-340.
- Cameotra, S.S. and Makkar, R.S. 1998. Synthesis of biosurfactants in extreme conditions. *Appl. Microbiol. Biotechnol.*, 50 : 520-529.
- Desai, J.D. and Banat, I.M. 1997. Microbial production of surfactants and their commercial potential. *Microbiol. Mol. Biol. Rev.*, 61 : 47-64.
- Djomo, J.E., Dauta, A., Ferrier, V., Narbonne, J.F., Monkiedje, A., Njine, T. and Garrigues, P. 2004. Toxic effects of some major polyaromatic hydrocarbons found in crud oil and aquatic sediments on *Scenedesmus subspicatus*. *Water Res.*, 38 : 1817-1821.
- El-Sayed, A.H.M.M., Mahmoud, W.M., Davis, E.M. and Coughlin, R.C. 1996. Growth of hydrocarbon-utilizing isolates in chemically defined media. *Int. Biodeter. Biodegr.*, 37 : 61-68.
- Garcia-Junco, M., Gomez-Lahoz, C., Niqui-Arroyo, J.L. and Ortega-Calvo, J.J. 2003. Biosurfactant- and biodegradation-enhanced partitioning of polycyclic aromatic hydrocarbons from nonaqueous-phase liquids. *Environ. Sci. Technol.*, 37 : 2988-2996.
- Georgiou, G., Lin, S.C. and Sharma, M.M. 1990. Surface active compounds from microorganisms. *Bio/Technology* 10 : 60-65.
- Gilewicz, M., Nadalig, N.T., Budzinski, H., Doumenq, P., Michotey, V. and Bertrand, J.C. 1997. Isolation and characterization of a marine bacterium capable of utilizing 2-methylphenanthrene. *Appl. Microbiol. Biotechnol.*, 48 : 528-533.

- Goes, A.P. and Sheppard, J.D. 1999. Effect of surfactants on α -amylase production in a solid substrate fermentation process. *J. Chem. Technol. Biotechnol.*, 74 : 709-712.
- Haferburg, D., Hommel, R., Claus, R. and Kleber, H.P. 1986. Extracellular microbial lipids as biosurfactants. *Adv. Biocjem. Eng. Biotechnol.*, 33 : 53-93.
- Harayama, S., Kishira, H., Kasai, Y. and Shutsubo, K. 1999. Petroleum biodegradation in marine environments. *J. Mol. Microbiol. Biotechnol.*, 1 : 63-70.
- Hua, Z., Chen, J., Lun, S. and Wang, X. 2003. Influence of biosurfactants produced by *Candida Antarctica* on surface properties of microorganism and biodegradation of *n*-alkanes. *Water Res.*, 37 : 4143-4150.
- Husain, D.R., Goutx, M., Acquaviva, M., Gilewicz, M. and Bertrand, J.-C. 1997. The effect of temperature on eicosane substrate uptake modes by a marine bacterium *Pseudomonas nautica* strain 617: relationship with the biochemical content of cells and supernatants. *World J. Microbiol. Biotechnol.* 13 : 587-590.
- Huy, N.Q., Jin, S.J., Amada, K., Haruki, M., Huu, N.B., Hang, D.T., Ha, D.T.C., Imanaka, T., Morikawa, M. and Kanaya, S. 1999. Characterization of petroleum-degrading bacteria from oil-contaminated sites in Vietnam. *J. Biosic. Bioeng.*, 88 : 100-102.
- Iwabuchi, N., Sunairi, M., Urai, M., Itoh, C., Anzai, H., Nakajima, M. and Harayama, S. 2002. Extracellular polysaccharides of *Rhodococcus rhodochrous* S-2 stimulate the degradation of aromatic components in crude oil by indigenous marine bacteria. *Appl. Environ. Microbiol.*, 68 : 2337-2343.
- Käppeli, O., Walther, P., Mueller, M. and Fiechter, A. 1984. Structure of the cell surface of the yeast *Candida tropicalis* and its relation to hydrocarbon transport. *Arch. Microbiol.*, 138 : 279-282.
- Kim, H.S, Yoon, B.D., Lee, C.H., Suh, H.H., Oh, H.M., Katsuragi, T. and Tani, Y. 1997. Production and properties of a lipopeptide biosurfactant from *Bacillus subtilis* C9. *J. Ferment. Bioeng.*, 84 : 41-46.
- Komukai-Nakamura, S., Sugiura, K., Yamauchi-Inomata, Y., Toki, H., Venkateswaran, K., Yamamoto, S., Tanaka, H. and harayama, S. 1996. Construction of bacterial consortia that degrade Arabian light crude oil. *J. Ferment. Bioeng.*, 82 : 570-574.

- Lee, S.C., Yoo, J.S., Kim, S.H., Chung, S.Y., Hwang, C.W., Joo, W.H. and Choi, Y.L. 2006. Production and characterization of lipopeptide biosurfactant from *Bacillus subtilis* A8-8. *J. Microbiol. Biotechnol.*, 16: 716-723.
- Lin, S.C., Lin, K.G., Lo, C.C. and Lin, Y.M. 1998. Enhanced biosurfactant production by a *Bacillus licheniformis* mutant. *Enzyme Microb. Technol.*, 23 : 267-273.
- Lin, T.C., Young, C.C., Ho, M.J., Yeh, M.S., Chou, C.L., Wei, Y.H. and Chang, J.S. 2005. Characterization of floating activity of indigenous diesel-assimilating bacterial isolates. *J. Biosci. Bioeng.*, 99 : 466-472.
- Luna-Velasco, M.A., Esparza-Garcia, F., Canizares-Villanueva, R.O. and Rodriguez-Vazquez, R. 2007. Production and properties of a bioemulsifier synthesized by phenanthrene-degrading *Penicillium* sp. *Process Biochem.*, 42 : 310-314.
- Maki, H., Hirayama, N., Hiwatari, T., Kohata, K., Uchiyama, H., Watanabe, M., Yamasaki, F. and Furuki, M. 2003. Crude oil bioremediation field experiment in the Sea of Japan. *Mar. Pollut. Bull.*, 47 : 74-77.
- Maneerat, S., Nitoda, T., Kanzaki, H. and Kawai, F. 2005. Bile acids are new products of a marine bacterium, *Myroides* sp. strain SM1. *Appl. Microbiol. Biotechnol.* 67 : 679-683.
- Maneerat, S., Bamba, T., Harada, K., Kobayashi, A., Yamada, H. and Kawai, F. 2006. A novel crude oil emulsifier excreted in the culture supernatant of a marine bacterium, *Myroides* sp. strain SM1. *Appl. Microbiol. Biotechnol.* 70 : 254-259.
- Marín, M., Pedregosa, A., Ríos, S., Ortiz, L. and Laborda, F. 1995. Biodegradation of diesel and heating oil by *Acinetobacter calcoaceticus* MM5: its possible applications on bioremediation. *Int. Biodeter. Biodegra.*, 35 : 269-285.
- McInerney, M.J., Javaheri, M. and Nagle Jr, D.P. 1990. Properties of the biosurfactant produced by *Bacillus licheniformis* strain JF-2. *J. Ind. Microbiol.*, 5 : 95-102.
- Mercadé, M.E., Monleón, L., de Andrés, C., Rodón, I., Martínez, E., Espuny, M.J. and Manresa, A. 1996. Screening and selection of surfactant-producing bacteria from waste lubricating oil. *J. Appl. Bacteriol.*, 81: 161-166.

- Meylheuc, T., van Oss, C.J. and Bellon-Fontaine, M.N. 2001. Adsorption of biosurfactant on solid surfaces and consequences regarding the bioadhesion of *Listeria monocytogenes* LO28. *J. Appl. Microbiol.*, 91: 822-832.
- Milewski, S. Protein Structure and Physicochemical Properties. **In:** Chemical and Functional Properties of Food Proteins. Sikorski, Z.E. (Ed) Technomic Publishing Company, Inc. Pennsylvania. pp.35-55.
- Morikawa, M., Ito, M. and Imanaka, T. 1992. Isolation of a new surfactin producer *Bacillus pumilus* A-1, and cloning and nucleotide sequence of the regulator gene, *psf-1*. *J. Ferment. Bioeng.*, 74: 255-261.
- Nakar, D. and Gutnick, D. 2001. Analysis of the *wee* gene cluster responsible for the biosynthesis of the polymeric bioemulsifier from the oil-degrading strain *Acinetobacter lwoffii* RAG-1. *Microbiology* 147 : 1937-1946.
- Nakata, K. 2000. Two glycolipids increase in the bioremediation of halogenated aromatic compounds. *J. Biosci. Bioeng.*, 89 : 577-581.
- Navon-Venezia, S., Zosim, Z., Gottlieb, A., Legmann, R., Carmeli, S., Ron, E.Z. and Rosenberg, E. 1995. Alasan, a new bioemulsifier from *Acinetobacter radioresistens*. *Appl. Environ. Microbiol.*, 61 : 3240-3244.
- Nitschke, M. and Pastore, G.M. 2006. Production and properties of a surfactant obtained from *Bacillus subtilis* grown on cassava wastewater. *Bioresource Technol.*, 97: 336-341.
- Olivera, N.L., Commendatore, M.G., Delgado, O. and Esteves, J.L. 2003. Microbial characterization and hydrocarbon biodegradation potential of natural bilge waste microflora. *J. Ind. Microbiol. Biotechnol.*, 30 : 542-548.
- Passeri, A., Schmidt, M., Haffner, T., Wray, V., Lang, S. and Wagner, F. 1992. Marine biosurfactants. IV. Production, characterization and biosynthesis of an anionic glucose lipid from the marine bacterial strain MM1. *Appl. Microbiol. Biotechnol.* 37 : 281-286.
- Pelletier, E., Delille, D. and Delille, B. 2004. Crude oil bioremediation in sub-Antarctic intertidal sediments: chemistry and toxicity of oiled residues. *Mar. Environ. Res.*, 57 : 311-327.

- Pini, F., Grossi, C., Nereo, S., Michaud, L., Giudice, A.L., Bruni, V., Baldi, F. and Fani, R. 2007. Molecular and physiological characterisation of psychrotrophic hydrocarbon-degrading bacteria isolated from Terra Nova Bay (Antarctica). *EUr. J. Soil Biol.* (in press).
- Poremba, K., Gunkel, W., Lang, S. and Wagner, F. 1991. Marine biosurfactants, III. Toxicity testing with marine microorganisms and comparison with synthetic surfactants. *Z. Naturforsch.* 46c : 210-216.
- Pruthi, V. and Cameotra, S.S. 1995. Rapid method for monitoring maximum biosurfactant production obtained by acetone precipitation. *Biotechnol. Techniques* 9 : 271-276.
- Radwan, S.S., Barabas, G., Sorkhoh, N.A., Damjanovich, S., Szabo, I., Szollosi, J., Matko, J., Penyige, A., Hirano, T. and Szabo, I.M. 1998. Hydrocarbon uptake by *Streptomyces*. *FEMS Microbiol. Lett.*, 169 : 87-94.
- Ron, E.Z. and Rosenberg, E. 2001. Natural roles of biosurfactants. *Environ. Microbiol.*, 3 : 229-236.
- Roongsawang, N., Thaniyavarn, J. and Thaniyavarn, S. 1999. Properties of biosurfactant produced by *Bacillus* sp. strain KP-2. *Thai J. Biotechnol.*, 1 : 54-60.
- Rosenberg, M., Gutnick, D. and Rosenberg, E. 1980. Adherence of bacteria to hydrocarbons: a simple method for measuring cell surface hydrophobicity. *FEMS Microbiol. Lett.* 9 : 29-33.
- Rosenberg, E. and Ron, E.Z. 1999. High- and low-molecular-mass microbial surfactants. *Appl. Microbiol. Biotechnol.*, 52 : 154-162.
- Schulz, D., Passeri, A., Schmidt, M., Lang, S., Wagner, F., Wray, V. and Gunkel, W. 1991. Marine biosurfactants, I. Screening for biosurfactants among crude oil degrading marine microorganisms from the North Sea. *Z. Naturforsch.* 46c : 197-203.
- Shriadah, M.M.A. 1998. Impacts of an oil spill on the marine environment of the United Arab Emirates along the Gulf of Oman. *Mar. Pollut. Bull.*, 36 : 876-879.
- Southam, G., Whitney, M. and Knickerbocker, C. 2001. Structural characterization of the hydrocarbon degrading bacteria-oil interface: implications for bioremediation. *Int. Biodeter. Biodegr.*, 47 : 197-201.
- Tuleve, B.K., Ivanov, G.R. and Christova, N.E. 2002. Biosurfactant production by a new *Pseudomonas putida* strain. *Z. Naturforsch.*, 57c : 356-360.

- Wilkinson, B.J., Sment, K.A. and Mayberry, W.R. 1982. Occurrence, localization, and possible significance of an ornithine-containing lipid in *Paracoccus denitrificans*. Arch. Microbiol., 131 : 338-343.
- Wilkinson, S.G. 1972. Composition and structure of the ornithine-containing lipid from *Pseudomonas rubescens*. Biochim. Biophys. Acta 270 : 1-17.
- Venkateswaran, K., Hoaki, T., Kato, M. and Maruyama, T. 1995. Microbial degradation of resins fractionated from Arabian light crude oil. Can. J. Microbiol., 41 : 418-424.
- Verdin, A., Sahraoui, A.L-H., Newsam, R., Robinsos, G. and Durand, R. 2005. Polycyclic aromatic hydrocarbons storage by *Fusarium solani* in intracellular lipid vesicles. Environ. Pollut., 133 : 283-291.
- Yakimov, M.M., Golyshin, P.N., Lang, S., Moore, E.R.B., Abraham, W.R., Lünsdorf, H. and Timmis, K.N. 1998. *Alcanivorax borkumensis* gen. nov. sp., nov., a new, hydrocarbon-degrading and surfactant-producing marine bacterium. Int. J. Syst. Bacteriol., 48 : 339-348.
- Yakimov, M.M., Timmis, K.N., Wray, V. and Fredrickson, H.L. 1995. Characterization of a new lipopeptide surfactant produced by thermotolerant and halotolerant subsurface *Bacillus licheniformis* BAS50. Appl. Environ. Microbiol., 61 : 1706-1713.
- Zajic, J.E., Gignard, H. and Gerson, D.F. 1977. Properties and biodegradation of a bioemulsifier from *Corynebacterium hydrocarboclastus*. Biotechnol. Bioeng., 19 : 1303-1320.
- Zinjarde, S.S. and Pant, A. 2002. Emulsifier from a tropical marine yeast, *Yarrowia lipolytica* NCIM 3589. J. Basic Microbiol. 42 : 67-73.